

GEOMETRIC APPROACH TO PARABOLIC INDUCTION

DAVID KAZHDAN AND YAKOV VARSHAVSKY

ABSTRACT. In this note we construct a “restriction” map from the cocenter of a reductive group G over a local non-archimedean field F to the cocenter of a Levi subgroup. We show that the dual map corresponds to parabolic induction and deduce that parabolic induction preserves stability. We also give a new (purely geometric) proof that the character of normalized parabolic induction does not depend on the parabolic subgroup. In the appendix, we use a similar argument to extend a theorem of Lusztig–Spaltenstein on induced unipotent classes to all infinite fields.

To Iosif Bernstein with gratitude and best wishes on his birthday

CONTENTS

Introduction	1
1. Preliminaries on algebraic groups	3
2. Top degree differential forms	8
3. Smooth measures, restriction, and induction	10
4. Relation to characters of induced representations	14
5. Restriction to an equal rank subgroup	16
6. Normalized induction, independence of $\mathbf{P}$, and stability	19
Appendix A. A generalization of a theorem of Lusztig–Spaltenstein	22
Appendix B. On a theorem of Harish–Chandra	26
References	29

INTRODUCTION

Let $\mathbf{G}$ be a linear algebraic group over a local non-archimedean field F , and let $G = \mathbf{G}(F)$. We denote by $\mathcal{H}(G)$ the space of smooth measures with compact support on G , by $\mathcal{H}(G)_G$ the space of the coinvariants of $\mathcal{H}(G)$ with respect to the

Date: November 26, 2015.

D.K. was supported by the ERC grant No. 247049-GLC, Y.V. was supported by the ISF grant 1017/13.

adjoint action of G , and let $\widehat{C}^G(G) := \text{Hom}_{\mathbb{C}}(\mathcal{H}(G)_G, \mathbb{C})$ be the space of invariant generalized functions on G . To every admissible representation π of G , one can associate its character $\chi_\pi \in \widehat{C}^G(G)$.

Now assume that $\mathbf{G}$ is connected reductive, $\mathbf{P} \subset \mathbf{G}$ a parabolic subgroup, $\mathbf{M} \subset \mathbf{P}$ a Levi subgroup, and $\mathbf{U} \subset \mathbf{P}$ the unipotent radical. For every admissible representation ρ of $M = \mathbf{M}(F)$ we denote by $\pi = i_{\mathbf{P}, \mathbf{M}}^G(\rho)$ the admissible representation of G obtained by the normalized parabolic induction of ρ .

In this work we construct a continuous map $i_{\mathbf{P}, \mathbf{M}}^G : \widehat{C}^M(M) \rightarrow \widehat{C}^G(G)$, satisfying $i_{\mathbf{P}, \mathbf{M}}^G(\chi_\rho) = \chi_\pi$ for every ρ as above. Namely, we construct a “restriction” map $r_{\mathbf{P}, \mathbf{M}}^G : \mathcal{H}(G)_G \rightarrow \mathcal{H}(M)_M$, and define $i_{\mathbf{P}, \mathbf{M}}^G$ to be the linear dual of $r_{\mathbf{P}, \mathbf{M}}^G$.

Then we show that $i_{\mathbf{P}, \mathbf{M}}^G$ does not depend on $\mathbf{P} \supset \mathbf{M}$. From this we conclude that for each ρ as above, the set of composition factors of $i_{\mathbf{P}, \mathbf{M}}^G(\rho)$ does not depend on $\mathbf{P}$. This gives a geometric proof of [BDK, Lem 5.4 (iii)]¹.

Finally, we show that $i_{\mathbf{P}, \mathbf{M}}^G$ preserves stability. This implies that parabolic induction of a stable representation is stable. This result is considered to be well-known by specialists, but does not seem to appear in a written form.

In the first appendix, we study a related question, motivated by the work of Lusztig–Spaltenstein [LS].

Let $\mathbf{G}$ and $\mathbf{P}$, $\mathbf{M}$ and $\mathbf{U}$ be as above, but over an arbitrary infinite field F . To every unipotent conjugacy class $C \subset M$ we associate an $\text{Ad } G$ -invariant subset $C_{P;G} := \cup_{g \in G} g(C \cdot U)g^{-1} \subset G$. Then $C_{P;G}$ is a union of unipotent conjugacy classes in G . A natural question is to what extent the set $C_{P;G}$ depends on $\mathbf{P} \supset \mathbf{M}$.

In their work, Lusztig and Spaltenstein [LS] showed, using representation theory, that the Zariski closure of $C_{P;G}$ does not depend on $\mathbf{P}$. A simpler proof of this fact was given later by Lusztig [Lu, Lem 10.3(a)]. This result can be thought of as the assertion that $C_{P;G}$ is “essentially” independent of $\mathbf{P}$, if F is algebraically closed.

We extend this result to an arbitrary F . Namely, for every algebraic variety $\mathbf{X}$ over F and a subset $A \subset X = \mathbf{X}(F)$ we define a “saturation” $\text{sat}(A) \subset X$. The main result of the appendix asserts that the saturation $\text{sat}(C_{P;G}) \subset G$ does not depend on $\mathbf{P}$.

Since every Zariski closed subset is saturated, our result is an extension of the theorem of Lusztig–Spaltenstein. Similarly, when F is a local field, our result implies that the closure of $C_{P;G}$ in the analytic topology does not depend on $\mathbf{P}$.

¹It was shown in [Ka, Thm B] that the span of characters of smooth irreducible representations of G is dense in $\widehat{C}^G(G)$. Therefore the independence assertion for $i_{\mathbf{P}, \mathbf{M}}^G$ follows from the corresponding result for characters ([BDK, Lem 5.4 (iii)]).

The result of the appendix indicates that the independence from P of normalized parabolic induction has a purely algebraic flavour. We also believe that the notion of saturation is interesting in its own right and deserves to be studied.

In the second appendix we give a proof of a group version of a theorem of Harish–Chandra, asserting that in characteristic zero the span of regular semisimple orbital integrals is dense in $\widehat{C}^G(G)$. Though this very important result is considered to be well-known among specialists, only its Lie algebra analog [HC, Thm 3.1] seems to appear in the literature. In particular, this fills a small gap in the argument of [Ka, Thm B].

Our paper is organized as follows. In Section 1 we describe general properties of reductive groups and so-called generalized Grothendieck–Springer resolutions. In Section 2 we study non-vanishing top degree differential forms, which are basic tools for this work. Note that in these two sections the ground field F is arbitrary, while starting from Section 3 the field F is local non-archimedean.

In Section 3 we introduce smooth measures with compact support and carry out our construction of the restriction $r_{P;M}^G$ and the induction $i_{P;M}^G$. In Section 4 we show that the induction map sends a character of a representation to a character of the induced representation. Then, in Section 5 we construct another restriction map R_H^G , defined for a connected equal rank subgroup $\mathbf{H}$ of $\mathbf{G}$, and show that R_H^G is compatible with $r_{P;M}^G$. Finally, in Section 6 we deduce from the results of Section 5 that the normalized parabolic induction is independent of $\mathbf{P}$ and preserves stability.

Notice that in Sections 1-5, we work with non-normalized induction, which has a purely geometric interpretation, while we pass to normalized induction only in Section 6.

We thank Yuval Flicker and the referee for a number of helpful suggestions and corrections.

1. PRELIMINARIES ON ALGEBRAIC GROUPS

1.1. The Chevalley map. (a) Let $\mathbf{G}$ be a linear algebraic group over a field F , let $\mathbf{c}_{\mathbf{G}} := \operatorname{Spec} F[\mathbf{G}]^{\mathbf{G}}$ be the Chevalley space of $\mathbf{G}$, and let $\nu_{\mathbf{G}} : \mathbf{G} \rightarrow \mathbf{c}_{\mathbf{G}}$ be the morphism dual to the inclusion $F[\mathbf{G}]^{\mathbf{G}} \hookrightarrow F[\mathbf{G}]$.

(b) A homomorphism of linear algebraic groups $\mathbf{H} \rightarrow \mathbf{G}$ induces a morphism $\pi_{\mathbf{H},\mathbf{G}} : \mathbf{c}_{\mathbf{H}} \rightarrow \mathbf{c}_{\mathbf{G}}$ of the Chevalley spaces, making the following diagram commutative

$$\begin{array}{ccc} \mathbf{H} & \xrightarrow{\nu_{\mathbf{H}}} & \mathbf{c}_{\mathbf{H}} \\ \downarrow & & \downarrow \pi_{\mathbf{H},\mathbf{G}} \\ \mathbf{G} & \xrightarrow{\nu_{\mathbf{G}}} & \mathbf{c}_{\mathbf{G}}. \end{array}$$

(c) Let $\mathbf{G}$ be connected reductive and split, $\mathbf{T} \subset \mathbf{G}$ a maximal split torus, and $W_{\mathbf{G}} = W_{\mathbf{G}, \mathbf{T}}$ the Weyl group of $\mathbf{G}$. Then the restriction $\nu_{\mathbf{G}}|_{\mathbf{T}} : \mathbf{T} \rightarrow \mathbf{c}_{\mathbf{G}}$ is surjective and induces an isomorphism $W_{\mathbf{G}} \backslash \mathbf{T} \xrightarrow{\sim} \mathbf{c}_{\mathbf{G}}$ (see [St, Cor 6.4]).

1.2. Notation. Let $\mathbf{G}$ be as in 1.1, and let $\mathbf{H} \subset \mathbf{G}$ be a closed subgroup.

(a) Let $\lambda_{\mathbf{G}} : \mathbf{G} \rightarrow \mathbb{G}_m$ be the homomorphism $g \mapsto \det \text{Ad } g$ and let $\Delta_{\mathbf{H}, \mathbf{G}} \in F[\mathbf{H}]^{\mathbf{H}} = F[\mathbf{c}_{\mathbf{H}}]$ be the $\text{Ad } \mathbf{H}$ -invariant function $h \mapsto \det(\text{Ad } h^{-1} - 1, \text{Lie } \mathbf{G} / \text{Lie } \mathbf{H})$.

(b) Let $\mathbf{H}^{\text{reg}/\mathbf{G}} \subset \mathbf{H}$ (resp. $\mathbf{c}_{\mathbf{H}}^{\text{reg}/\mathbf{G}} \subset \mathbf{c}_{\mathbf{H}}$) be the open subscheme defined by the condition $\Delta_{\mathbf{H}, \mathbf{G}} \neq 0$. By construction, $\mathbf{H}^{\text{reg}/\mathbf{G}} = \nu_{\mathbf{H}}^{-1}(\mathbf{c}_{\mathbf{H}}^{\text{reg}/\mathbf{G}})$.

(c) Let $\mathbf{K} \subset \mathbf{H}$ be a closed subgroup. Then $\Delta_{\mathbf{K}, \mathbf{G}} = \Delta_{\mathbf{K}, \mathbf{H}} \cdot (\Delta_{\mathbf{H}, \mathbf{G}}|_{\mathbf{K}})$, thus $\mathbf{K}^{\text{reg}/\mathbf{G}} = \mathbf{K}^{\text{reg}/\mathbf{H}} \cap \mathbf{H}^{\text{reg}/\mathbf{G}}$.

1.3. Remark. Notice that subset $\mathbf{H}^{\text{reg}/\mathbf{G}} \subset \mathbf{H}$ should not be confused with a more standard subset $\mathbf{H}^{\mathbf{G}-\text{reg}} := \mathbf{H} \cap \mathbf{G}^{\text{reg}} \subset \mathbf{H}$.

1.4. The generalized Grothendieck–Springer resolution.

(a) Let $\mathbf{H}$ (see 1.2) act on an algebraic variety $\mathbf{X}$ over F . We denote by $\text{Ind}_{\mathbf{H}}^{\mathbf{G}}(\mathbf{X}) := \mathbf{G} \times^{\mathbf{H}} \mathbf{X}$ the quotient of $\mathbf{G} \times \mathbf{X}$ by $\mathbf{H}$, acting by $h(g, x) := (gh^{-1}, h(x))$. Then $\text{Ind}_{\mathbf{H}}^{\mathbf{G}}(\mathbf{X})$ is equipped with an action of $\mathbf{G}$, given by $g'([g, h]) = [g'g, h]$.

(b) Assume that $\mathbf{H}$ acts on itself by conjugation, and set $\tilde{\mathbf{G}}_{\mathbf{H}} := \text{Ind}_{\mathbf{H}}^{\mathbf{G}}(\mathbf{H})$. Explicitly, $\tilde{\mathbf{G}}_{\mathbf{H}} = \mathbf{G} \times^{\mathbf{H}, \text{Ad}} \mathbf{H}$, where Ad indicates the adjoint action.

(c) We have a natural closed embedding $(\text{pr}_{\mathbf{G}/\mathbf{H}}, a_{\mathbf{H}, \mathbf{G}}) : \tilde{\mathbf{G}}_{\mathbf{H}} \hookrightarrow (\mathbf{G}/\mathbf{H}) \times \mathbf{G}$, defined by $[g, h] \mapsto ([g], ghg^{-1})$, whose image consists of all pairs $([g], x)$ such that for every representative $g \in \mathbf{G}$ of $[g]$ we have $x \in \mathbf{H}_g := g\mathbf{H}g^{-1}$. In particular, the projection $\text{pr}_{\mathbf{G}/\mathbf{H}} : \tilde{\mathbf{G}}_{\mathbf{H}} \rightarrow \mathbf{G}/\mathbf{H}$ is smooth, and for every $[g] \in \mathbf{G}/\mathbf{H}$, the map $a_{\mathbf{H}, \mathbf{G}} : \tilde{\mathbf{G}}_{\mathbf{H}} \rightarrow \mathbf{G}$ identifies the fiber $\text{pr}_{\mathbf{G}/\mathbf{H}}^{-1}([g])$ with $\mathbf{H}_g$.

1.5. Simple properties. (a) Every $\text{Ad } \mathbf{H}$ -invariant morphism $f : \mathbf{H} \rightarrow \mathbf{X}$ gives rise to an $\text{Ad } \mathbf{G}$ -equivariant morphism $\tilde{\mathbf{G}}_{\mathbf{H}} \rightarrow \mathbf{X} : [g, h] \mapsto f(h)$. In particular, $\lambda_{\mathbf{H}}$ gives rise to a morphism $\lambda_{\tilde{\mathbf{G}}_{\mathbf{H}}} : \tilde{\mathbf{G}}_{\mathbf{H}} \rightarrow \mathbb{G}_m$, and $\Delta_{\mathbf{H}, \mathbf{G}}$ gives rise to a function $\Delta_{\tilde{\mathbf{G}}_{\mathbf{H}}} : \tilde{\mathbf{G}}_{\mathbf{H}} \rightarrow \mathbb{A}^1$.

(b) By (a), the morphism $\nu_{\mathbf{H}} : \mathbf{H} \rightarrow \mathbf{c}_{\mathbf{H}}$ gives rise to a morphism $\nu_{\tilde{\mathbf{G}}_{\mathbf{H}}} : \tilde{\mathbf{G}}_{\mathbf{H}} \rightarrow \mathbf{c}_{\mathbf{H}}$. Moreover, by 1.1(b), we have a commutative diagram

$$\begin{array}{ccc} \tilde{\mathbf{G}}_{\mathbf{H}} & \xrightarrow{\nu_{\tilde{\mathbf{G}}_{\mathbf{H}}}} & \mathbf{c}_{\mathbf{H}} \\ a_{\mathbf{H}, \mathbf{G}} \downarrow & & \downarrow \pi_{\mathbf{H}, \mathbf{G}} \\ \mathbf{G} & \xrightarrow{\nu_{\mathbf{G}}} & \mathbf{c}_{\mathbf{G}}, \end{array}$$

which induces a morphism $\tilde{\mathbf{G}}_{\mathbf{H}} \rightarrow \mathbf{G} \times_{c_{\mathbf{G}}} c_{\mathbf{H}}$.

(c) Set $\tilde{\mathbf{G}}_{\mathbf{H}}^{\text{reg}} := \nu_{\tilde{\mathbf{G}}_{\mathbf{H}}}^{-1}(c_{\mathbf{H}}^{\text{reg}/\mathbf{G}}) \subset \tilde{\mathbf{G}}_{\mathbf{H}}$. Explicitly, $\tilde{\mathbf{G}}_{\mathbf{H}}^{\text{reg}} = \text{Ind}_{\mathbf{H}}^{\mathbf{G}}(\mathbf{H}^{\text{reg}/\mathbf{G}})$. By definition, the morphism of (b) induces a morphism $\iota_{\mathbf{H},\mathbf{G}} : \tilde{\mathbf{G}}_{\mathbf{H}}^{\text{reg}} \rightarrow \mathbf{G} \times_{c_{\mathbf{G}}} c_{\mathbf{H}}^{\text{reg}/\mathbf{G}}$.

(d) Let $\mathbf{K} \subset \mathbf{H}$ be a closed subgroup. Then the morphism $a_{\mathbf{K},\mathbf{H}} : \tilde{\mathbf{H}}_{\mathbf{K}} \rightarrow \mathbf{H}$ induces a morphism $\tilde{\mathbf{G}}_{\mathbf{K}} = \text{Ind}_{\mathbf{H}}^{\mathbf{G}}(\tilde{\mathbf{H}}_{\mathbf{K}}) \rightarrow \text{Ind}_{\mathbf{H}}^{\mathbf{G}}(\mathbf{H}) = \tilde{\mathbf{G}}_{\mathbf{H}}$, which we again denote by $a_{\mathbf{K},\mathbf{H}}$. Note that the composition $a_{\mathbf{H},\mathbf{G}} \circ a_{\mathbf{K},\mathbf{H}} : \tilde{\mathbf{G}}_{\mathbf{K}} \rightarrow \tilde{\mathbf{G}}_{\mathbf{H}} \rightarrow \mathbf{G}$ is equal to $a_{\mathbf{K},\mathbf{G}}$. By 1.2(c), $a_{\mathbf{K},\mathbf{H}}$ induces a morphism $\tilde{\mathbf{G}}_{\mathbf{K}}^{\text{reg}} \rightarrow \tilde{\mathbf{G}}_{\mathbf{H}}^{\text{reg}}$.

(e) We also set $\tilde{\mathbf{H}}_{\mathbf{K}}^{\text{reg}/\mathbf{G}} := \nu_{\tilde{\mathbf{H}}_{\mathbf{K}}}^{-1}(c_{\mathbf{K}}^{\text{reg}/\mathbf{G}}) \subset \tilde{\mathbf{H}}_{\mathbf{K}}$.

1.6. The equal rank case. (a) Let $\mathbf{G}$ be connected reductive, and let $\mathbf{H} \subset \mathbf{G}$ be a connected *equal rank* subgroup, by which we mean that a maximal torus $\mathbf{T} \subset \mathbf{H}$ is a maximal torus of $\mathbf{G}$. Let $\mathbf{U} = \mathbf{U}_{\mathbf{H}} \subset \mathbf{H}$ be the unipotent radical of $\mathbf{H}$, and fix a maximal torus $\mathbf{T} \subset \mathbf{H}$, defined over F .

(b) Assume that $\mathbf{T}$ is split. Then, by definition, an element $t \in \mathbf{T}$ belongs to $\mathbf{H}^{\text{reg}/\mathbf{G}}$ if and only if $\alpha(t) \neq 0$ for every root $\alpha \in \Phi(\mathbf{G}, \mathbf{T}) \setminus \Phi(\mathbf{H}, \mathbf{T})$. Equivalently, this happens if and only if the connected stabilizer $\mathbf{Z}_{\mathbf{G}}(t)^0$ equals $\mathbf{Z}_{\mathbf{H}}(t)^0$. If, in addition, the derived group of $\mathbf{G}$ is simply connected, then the stabilizer $\mathbf{Z}_{\mathbf{G}}(t)$ is connected. In this case, for every $t \in \mathbf{T} \cap \mathbf{H}^{\text{reg}/\mathbf{G}}$, we have $\mathbf{Z}_{\mathbf{G}}(t) = \mathbf{Z}_{\mathbf{H}}(t) \subset \mathbf{H}$.

Claim 1.7. *In the situation 1.6 (a), the group $\mathbf{H}$ has a Levi subgroup (see [Bo, 11.22]). Moreover, there exists a unique Levi subgroup $\mathbf{M} \subset \mathbf{H}$, containing $\mathbf{T}$.*

Proof. By uniqueness, we can extend scalars to a finite separable extension, thus assuming that $\mathbf{T}$ is split. In this case, the subgroup $\mathbf{H} \subset \mathbf{G}$ is generated by $\mathbf{T}$ and the root subgroups $\mathbf{U}_{\alpha} \subset \mathbf{G}$ for $\alpha \in \Phi(\mathbf{H}, \mathbf{T})$. Indeed, this follows from the fact that $\mathbf{H}$ is generated by its Borel subgroups $\mathbf{B} \supset \mathbf{T}$, and that the corresponding assertion for solvable $\mathbf{H}$ follows from [Bo, Prop 14.4].

Next we observe that the set of roots $\Phi(\mathbf{U}, \mathbf{T})$ consists of all $\alpha \in \Phi(\mathbf{H}, \mathbf{T})$ such that $-\alpha \notin \Phi(\mathbf{H}, \mathbf{T})$. Therefore a Levi subgroup $\mathbf{M} \supset \mathbf{T}$ of $\mathbf{H}$ has to coincide with the subgroup $\mathbf{M}' \subset \mathbf{G}$ generated by $\mathbf{T}$ and the roots subgroups $\mathbf{U}_{\alpha} \subset \mathbf{G}$ for all $\alpha \in \Phi' := \Phi(\mathbf{H}, \mathbf{T}) \setminus \Phi(\mathbf{U}, \mathbf{T})$.

It remains to show that the subgroup $\mathbf{M}' \subset \mathbf{H}$ defined in the previous paragraph is indeed a Levi subgroup. Since $\mathbf{M}'\mathbf{U} = \mathbf{H}$, it suffices to show that $\Phi(\mathbf{M}', \mathbf{T}) \subset \Phi'$. Recall that $\mathbf{G}$ has an automorphism ι such that $\iota(\mathbf{T}) = \mathbf{T}$ and $\iota(\mathbf{U}_{\alpha}) = \mathbf{U}_{-\alpha}$ for every $\alpha \in \Phi(\mathbf{G}, \mathbf{T})$. Since the subset $\Phi' \subset \Phi(\mathbf{G}, \mathbf{T})$ is stable under the map $\alpha \mapsto -\alpha$, we conclude that $\mathbf{M}' \subset \mathbf{H} \cap \iota(\mathbf{H})$, thus $\Phi(\mathbf{M}', \mathbf{T}) \subset \Phi(\mathbf{H}, \mathbf{T}) \cap \Phi(\iota(\mathbf{H}), \mathbf{T}) = \Phi'$. $\square$

1.8. Remark. In this work we only consider the case when $\mathbf{H}$ is either a parabolic subgroup of $\mathbf{G}$ or a connected centralizer of a semisimple element. In these cases,

Claim 1.7 is well-known (see [Bo, Cor 14.19]). On the other hand, we believe that the context of equal rank subgroups is the “correct framework” to work in.

Lemma 1.9. *In the notation of Claim 1.7, let p be the projection $\mathbf{H} \rightarrow \mathbf{H}/\mathbf{U} \cong \mathbf{M}$, and let i be the inclusion $\mathbf{M} \hookrightarrow \mathbf{H}$.*

(a) *Set $\mathbf{H}' := p^{-1}(\mathbf{M}^{\text{reg}/\mathbf{G}}) \subset \mathbf{H}$. Then the morphism $\text{Ind}_{\mathbf{M}}^{\mathbf{H}}(\mathbf{M}^{\text{reg}/\mathbf{G}}) \rightarrow \mathbf{H}'$, induced by $a_{\mathbf{M},\mathbf{H}} : \tilde{\mathbf{H}}_{\mathbf{M}} \rightarrow \mathbf{H}$, is an isomorphism.*

(b) *The morphisms $\pi_{\mathbf{H},\mathbf{M}} : \mathbf{c}_{\mathbf{H}} \rightarrow \mathbf{c}_{\mathbf{M}}$ and $\pi_{\mathbf{M},\mathbf{H}} : \mathbf{c}_{\mathbf{M}} \rightarrow \mathbf{c}_{\mathbf{H}}$, induced by p and i respectively (see 1.1(b)), are isomorphisms.*

(c) *The isomorphism $\pi_{\mathbf{M},\mathbf{H}} : \mathbf{c}_{\mathbf{M}} \xrightarrow{\sim} \mathbf{c}_{\mathbf{H}}$ induces an isomorphism $\mathbf{c}_{\mathbf{M}}^{\text{reg}/\mathbf{G}} \xrightarrow{\sim} \mathbf{c}_{\mathbf{H}}^{\text{reg}/\mathbf{G}}$.*

(d) *We have the equality $p^{-1}(\mathbf{M}^{\text{reg}/\mathbf{G}}) = \mathbf{H}^{\text{reg}/\mathbf{G}}$.*

(e) *The morphisms $\text{Ind}_{\mathbf{M}}^{\mathbf{H}}(\mathbf{M}^{\text{reg}/\mathbf{G}}) \rightarrow \mathbf{H}^{\text{reg}/\mathbf{G}}$ and $\tilde{\mathbf{G}}_{\mathbf{M}}^{\text{reg}} \rightarrow \tilde{\mathbf{G}}_{\mathbf{H}}^{\text{reg}}$ induced by $a_{\mathbf{M},\mathbf{H}}$ (compare 1.5(d)) are isomorphisms.*

(f) *Set $e_{\mathbf{G}} := \nu_{\mathbf{G}}(1) \in \mathbf{c}_{\mathbf{G}}$ and similarly for $\mathbf{H}$. Then the morphism $\pi_{\mathbf{H},\mathbf{G}} : \mathbf{c}_{\mathbf{H}} \rightarrow \mathbf{c}_{\mathbf{G}}$ satisfies $\pi_{\mathbf{H},\mathbf{G}}^{-1}(e_{\mathbf{G}}) = e_{\mathbf{H}}$.*

Proof. (a) Since $\mathbf{H} = \mathbf{U} \rtimes \mathbf{M}$, it suffices to show that for every $m \in \mathbf{M}^{\text{reg}/\mathbf{G}}$ the map $f_m : \mathbf{U} \rightarrow \mathbf{U} : u \mapsto m^{-1}umu^{-1}$ is an isomorphism.

Consider the upper (or lower) filtration $\mathbf{U}^{(i)}$ of $\mathbf{U}$. It remains to show that f_m induces an isomorphism on each quotient $\mathbf{U}^{(i)}/\mathbf{U}^{(i+1)}$. Using the natural isomorphism $\mathbf{U}^{(i)}/\mathbf{U}^{(i+1)} \cong \text{Lie } \mathbf{U}^{(i)}/\mathbf{U}^{(i+1)}$, it suffices to show that $\text{Ad}(m^{-1}) - \text{Id}$ is invertible on $\text{Lie } \mathbf{U}^{(i)}/\mathbf{U}^{(i+1)}$. Since $m \in \mathbf{M}^{\text{reg}/\mathbf{G}}$, the map $\text{Ad}(m^{-1}) - \text{Id}$ is invertible on $\text{Lie } \mathbf{G}/\text{Lie } \mathbf{M}$. Hence it is invertible on $\text{Lie } \mathbf{U} = \text{Lie } \mathbf{H}/\text{Lie } \mathbf{M}$, and the assertion follows.

(b) Since $p \circ i = \text{Id}_{\mathbf{M}}$, it suffices to show that the pullback $i^* : F[\mathbf{H}]^{\mathbf{H}} \rightarrow F[\mathbf{M}]^{\mathbf{M}}$ is injective. By (a), the induced map $i^* : F[\mathbf{H}']^{\mathbf{H}} \rightarrow F[\mathbf{M}^{\text{reg}/\mathbf{G}}]^{\mathbf{M}}$ is injective. Since $\mathbf{H}' \subset \mathbf{H}$ is Zariski dense, the restriction map $F[\mathbf{H}] \rightarrow F[\mathbf{H}']$ is injective, and the assertion follows.

(c) Extending scalars to a finite separable extension, we can assume that $\mathbf{T}$ is split. Then, by 1.1(c), it remains to show that $\mathbf{T} \cap \mathbf{M}^{\text{reg}/\mathbf{G}} = \mathbf{T} \cap \mathbf{H}^{\text{reg}/\mathbf{G}}$. But this follows from the explicit description of both sides, given in 1.6(b).

The assertion (d) follows from (c), while (e) follows from (a) and (d).

(f) By (b), we can replace $\mathbf{H}$ by $\mathbf{M}$, thus assuming that $\mathbf{H}$ is reductive. Extending scalars, we can assume that $\mathbf{T}$ is split. In this case, $\pi_{\mathbf{H},\mathbf{G}} : \mathbf{c}_{\mathbf{H}} \rightarrow \mathbf{c}_{\mathbf{G}}$ is the projection $W_{\mathbf{H}} \setminus \mathbf{T} \rightarrow W_{\mathbf{G}} \setminus \mathbf{T}$ (see 1.1(c)), and the assertion is immediate. $\square$

Lemma 1.10. *In the situation of 1.6(a), assume that the derived group of $\mathbf{G}$ is simply connected. Then*

(a) *the morphism $\pi_{\mathbf{H},\mathbf{G}} : \mathbf{c}_{\mathbf{H}}^{\text{reg}/\mathbf{G}} \rightarrow \mathbf{c}_{\mathbf{G}}$ is étale.*

(b) *the morphism $\iota_{\mathbf{H},\mathbf{G}} : \tilde{\mathbf{G}}_{\mathbf{H}}^{\text{reg}} \rightarrow \mathbf{G} \times_{\mathbf{c}_{\mathbf{G}}} \mathbf{c}_{\mathbf{H}}^{\text{reg}/\mathbf{G}}$ from 1.5(c) is an isomorphism.*

Proof. Extending scalars to a finite separable extension of F , we can assume that $\mathbf{T}$ is split.

(a) By Lemma 1.9(c), we can replace $\mathbf{H}$ by its Levi subgroup $\mathbf{M}$, thus we can assume that $\mathbf{H}$ is reductive. Then $\pi_{\mathbf{H},\mathbf{G}} : \mathbf{c}_{\mathbf{H}} \rightarrow \mathbf{c}_{\mathbf{G}}$ is simply the projection $W_{\mathbf{H}} \backslash \mathbf{T} \rightarrow W_{\mathbf{G}} \backslash \mathbf{T}$ (see 1.1(c)). Thus, it remains to show that for every $t \in \mathbf{T} \cap \mathbf{H}^{\text{reg}}/\mathbf{G}$ we have the equality of stabilizers $\text{Stab}_{W_{\mathbf{H}}}(t) = \text{Stab}_{W_{\mathbf{G}}}(t)$. But our assumption on $\mathbf{G}$ implies that $\mathbf{Z}_{\mathbf{H}}(t) = \mathbf{Z}_{\mathbf{G}}(t)$ (see 1.6(b)), so the assertion follows.

(b) Note that $\tilde{\mathbf{G}}_{\mathbf{H}}^{\text{reg}}$ is étale over $\mathbf{G}$ by Corollary 2.6 below, while $\mathbf{G} \times_{\mathbf{c}_{\mathbf{G}}} \mathbf{c}_{\mathbf{H}}^{\text{reg}}/\mathbf{G}$ is étale over $\mathbf{G}$ by (a). Hence $\iota_{\mathbf{H},\mathbf{G}}$ is étale. Thus, in order to show that $\iota_{\mathbf{H},\mathbf{G}}$ is an isomorphism, it suffices to show that $\iota_{\mathbf{H},\mathbf{G}}$ is a bijection (on $\overline{F}$ -points).

First we show that $\iota_{\mathbf{H},\mathbf{G}}$ is surjective. Since $\nu_{\mathbf{H}}|_{\mathbf{T}} : \mathbf{T} \rightarrow \mathbf{c}_{\mathbf{H}}$ is surjective, every element of $\mathbf{G} \times_{\mathbf{c}_{\mathbf{G}}} \mathbf{c}_{\mathbf{H}}^{\text{reg}}/\mathbf{G}$ has a form $(g, \nu_{\mathbf{H}}(t))$ for some $g \in \mathbf{G}$ and $t \in \mathbf{T} \cap \mathbf{H}^{\text{reg}}/\mathbf{G}$ such that $\nu_{\mathbf{G}}(g) = \nu_{\mathbf{G}}(t)$. Let $g = su$ be the Jordan decomposition. Then $\nu_{\mathbf{G}}(s) = \nu_{\mathbf{G}}(g) = \nu_{\mathbf{G}}(t)$ (see [St, Cor 6.5]). Since s and t are semisimple, they are $\mathbf{G}$ -conjugate (by [St, Cor 6.6]). Since $\iota_{\mathbf{G},\mathbf{H}}$ is $\mathbf{G}$ -equivariant, we can replace g by its conjugate, thus assuming that $s = t$.

Since u is unipotent and $us = su$, we get that $u \in \mathbf{Z}_{\mathbf{G}}(s)^0$. Hence $g \in \mathbf{H}$ (use 1.6(b)), thus $g = su \in \mathbf{H}^{\text{reg}}/\mathbf{G}$, and $\iota_{\mathbf{H},\mathbf{G}}([1, g]) = (g, \nu_{\mathbf{H}}(t))$.

To show the injectivity, assume that two elements $\tilde{g} = [g, h]$ and $\tilde{g}' = [g', h']$ of $\tilde{\mathbf{G}}_{\mathbf{H}}^{\text{reg}}$ satisfy $\iota_{\mathbf{H},\mathbf{G}}(\tilde{g}) = \iota_{\mathbf{H},\mathbf{G}}(\tilde{g}')$. Since $\iota_{\mathbf{G},\mathbf{H}}$ is $\mathbf{G}$ -equivariant, we can replace $\tilde{g}$ and $\tilde{g}'$ by their g'^{-1} -conjugates, thus assuming that $g' = 1$. In this case, the identity $\iota_{\mathbf{H},\mathbf{G}}(\tilde{g}) = \iota_{\mathbf{H},\mathbf{G}}(\tilde{g}')$ implies that $\nu_{\mathbf{H}}(h) = \nu_{\mathbf{H}}(h')$ and $ghg^{-1} = h'$. It suffices to show that $g \in \mathbf{H}$, hence $\tilde{g} = [1, ghg^{-1}] = \tilde{g}'$.

Let $h = su$ and $h' = s'u'$ be the Jordan decompositions. Then $gsg^{-1} = s'$ and $\nu_{\mathbf{H}}(s) = \nu_{\mathbf{H}}(s')$. Thus, s and s' are $\mathbf{H}$ -conjugates. Hence, replacing (g, h) by (gx^{-1}, xhx^{-1}) for some $x \in \mathbf{H}$, we may assume that $s = s'$, thus $g \in \mathbf{Z}_{\mathbf{G}}(s)$. Since $h \in \mathbf{H}^{\text{reg}}/\mathbf{G}$, we get $s \in \mathbf{H}^{\text{reg}}/\mathbf{G}$. By 1.6(b), we conclude that $\mathbf{Z}_{\mathbf{G}}(s) \subset \mathbf{H}$, thus $g \in \mathbf{H}$. $\square$

Corollary 1.11. *In the situation of 1.6, the morphism $\iota_{\mathbf{H},\mathbf{G}} : \tilde{\mathbf{G}}_{\mathbf{H}}^{\text{reg}} \rightarrow \mathbf{G} \times_{\mathbf{c}_{\mathbf{G}}} \mathbf{c}_{\mathbf{H}}^{\text{reg}}/\mathbf{G}$ from 1.5(c) is finite and surjective.*

Proof. Let $\mathbf{G}^{\text{sc}}$ be the simply connected covering of the derived group of $\mathbf{G}$. Consider the natural isogeny $\pi : \mathbf{G}' := \mathbf{G}^{\text{sc}} \times \mathbf{Z}(\mathbf{G})^0 \rightarrow \mathbf{G}$, and set $\mathbf{H}' := \pi^{-1}(\mathbf{H}) \subset \mathbf{G}'$. Then we have a commutative diagram

$$\begin{array}{ccc} \tilde{\mathbf{G}}_{\mathbf{H}'}^{\text{reg}} & \xrightarrow{\iota_{\mathbf{H}',\mathbf{G}'}} & \mathbf{G}' \times_{\mathbf{c}_{\mathbf{G}'}} \mathbf{c}_{\mathbf{H}'}^{\text{reg}}/\mathbf{G}' \\ \downarrow & & \downarrow \\ \tilde{\mathbf{G}}_{\mathbf{H}}^{\text{reg}} & \xrightarrow{\iota_{\mathbf{H},\mathbf{G}}} & \mathbf{G} \times_{\mathbf{c}_{\mathbf{G}}} \mathbf{c}_{\mathbf{H}}^{\text{reg}}/\mathbf{G}, \end{array}$$

where vertical arrows are finite surjective morphisms, induced by π . Now, $\iota_{\mathbf{H}', \mathbf{G}'}$ is an isomorphism by Lemma 1.10. Therefore $\iota_{\mathbf{H}, \mathbf{G}}$ is finite and surjective. $\square$

1.12. Remark. Although the morphism $\iota_{\mathbf{H}, \mathbf{G}}$ from Corollary 1.11 is not an isomorphism in general, it is an isomorphism over a “strongly regular locus”. Moreover, the whole morphism $\iota_{\mathbf{H}, \mathbf{G}}$ “can be made an isomorphism”, if one replaces the (singular) Chevalley spaces $\mathbf{c}_{\mathbf{G}}$ and $\mathbf{c}_{\mathbf{H}}$ by their smooth Artin stack versions.

2. TOP DEGREE DIFFERENTIAL FORMS

2.1. Notation. (a) For every smooth algebraic (or analytic) variety $\mathbf{X}$ over F of dimension d , we denote by $\mathcal{K}_{\mathbf{X}} = \Omega_{\mathbf{X}}^d$ the canonical bundle on $\mathbf{X}$, that is, the line bundle of top degree differential forms on $\mathbf{X}$.

(b) For every map of smooth algebraic (or analytic) varieties $f : \mathbf{X} \rightarrow \mathbf{Y}$ over F of dimension d , the pullback map of differential forms gives rise to a morphism of line bundles $i_f : f^*(\mathcal{K}_{\mathbf{Y}}) \rightarrow \mathcal{K}_{\mathbf{X}}$.

2.2. The group case. Let $\mathbf{G}$ be an algebraic group over F .

(a) For each $g \in \mathbf{G}$, the left multiplication map $L_g : \mathbf{G} \rightarrow \mathbf{G} : x \mapsto gx$ induces an isomorphism $\text{Lie } \mathbf{G} = T_1(\mathbf{G}) \xrightarrow{\sim} T_g(\mathbf{G})$ of tangent spaces. These isomorphisms for all g induce a trivialization $\mathbf{G} \times \text{Lie } \mathbf{G} \xrightarrow{\sim} T(\mathbf{G})$ of the tangent bundle of $\mathbf{G}$.

(b) Consider the one-dimensional vector space $V_{\mathbf{G}} := \det(\text{Lie } \mathbf{G})^*$ over F . Then the trivialization from (a) induces an isomorphism $i : \mathcal{O}_{\mathbf{G}} \otimes_F V_{\mathbf{G}} \xrightarrow{\sim} \mathcal{K}_{\mathbf{G}}$ of line bundles. Explicitly, for every $v \in V_{\mathbf{G}}$ the differential form $i(v)$ is the unique left invariant differential form $\omega_{\mathbf{G}}(v) = \omega_{\mathbf{G}}^l(v)$ on $\mathbf{G}$ such that $\omega_{\mathbf{G}}^l(v)|_{g=1} = v$. Moreover, $\omega_{\mathbf{G}}^l(v)$ is nonvanishing if $v \neq 0$.

(c) Similarly, for every $v \in V_{\mathbf{G}}$ there exists a unique right invariant differential form $\omega_{\mathbf{G}}^r(v)$ on $\mathbf{G}$ such that $\omega_{\mathbf{G}}^r(v)|_{g=1} = v$. Explicitly, $\omega_{\mathbf{G}}^r(v) = \lambda_{\mathbf{G}} \cdot \omega_{\mathbf{G}}^l(v)$, where $\lambda_{\mathbf{G}}$ was defined in 1.2(a). Observe that if $\mathbf{G}$ is connected reductive, then the character $\lambda_{\mathbf{G}}$ is trivial, thus $\omega_{\mathbf{G}}^r(v) = \omega_{\mathbf{G}}^l(v)$.

2.3. The canonical bundle on $\tilde{\mathbf{G}}_{\mathbf{H}}$.

(a) For every $[g] \in \mathbf{G}/\mathbf{H}$ we have natural identifications $T_{[g]}(\mathbf{G}/\mathbf{H}) \xrightarrow{\sim} \text{Lie } \mathbf{G} / \text{Lie } \mathbf{H}_g$, and $\text{pr}_{\mathbf{G}/\mathbf{H}}^{-1}([g]) \xrightarrow{\sim} \mathbf{H}_g : [g, h] \mapsto ghg^{-1}$ (see 1.4(c)). Thus, for every $[g, h] \in \tilde{\mathbf{G}}_{\mathbf{H}}$, we have an exact sequence

$$0 \rightarrow T_{ghg^{-1}}(\mathbf{H}_g) \rightarrow T_{[g, h]}(\tilde{\mathbf{G}}_{\mathbf{H}}) \rightarrow \text{Lie } \mathbf{G} / \text{Lie } \mathbf{H}_g \rightarrow 0.$$

(b) Using the identification $\text{Lie } \mathbf{H}_g \xrightarrow{\sim} T_{ghg^{-1}}(\mathbf{H}_g)$, induced by $L_{ghg^{-1}}$ as in 2.2(a), we have a natural isomorphism

$$\det T_{[g, h]}(\tilde{\mathbf{G}}_{\mathbf{H}}) \cong \det \text{Lie } \mathbf{H}_g \otimes \det(\text{Lie } \mathbf{G} / \text{Lie } \mathbf{H}_g) \cong \det \text{Lie } \mathbf{G}.$$

(c) By (b), we have a natural isomorphism $i : \mathcal{O}_{\tilde{\mathbf{G}}_{\mathbf{H}}} \otimes_F V_{\mathbf{G}} \xrightarrow{\sim} \mathcal{K}_{\tilde{\mathbf{G}}_{\mathbf{H}}}$. In particular, every $0 \neq v \in V_{\mathbf{G}}$ defines a nonvanishing differential form $\omega_{\tilde{\mathbf{G}}_{\mathbf{H}}}(v) := i(v)$ on $\tilde{\mathbf{G}}_{\mathbf{H}}$.

(d) Let $a_{\mathbf{H},\mathbf{G}} : \tilde{\mathbf{G}}_{\mathbf{H}} \rightarrow \mathbf{G}$ be as in 1.4(c). Using (c) and the isomorphism $\mathcal{K}_{\mathbf{G}} \xrightarrow{\sim} \mathcal{O}_{\mathbf{G}} \otimes_F V_{\mathbf{G}}$ from 2.2(b), we get a natural isomorphism

$$\tilde{i} : a_{\mathbf{H},\mathbf{G}}^*(\mathcal{K}_{\mathbf{G}}) \xrightarrow{\sim} a_{\mathbf{H},\mathbf{G}}^*(\mathcal{O}_{\mathbf{G}} \otimes_F V_{\mathbf{G}}) = \mathcal{O}_{\tilde{\mathbf{G}}_{\mathbf{H}}} \otimes V_{\mathbf{G}} \xrightarrow{\sim} \mathcal{K}_{\tilde{\mathbf{G}}_{\mathbf{H}}}.$$

2.4. Remarks. (a) Recall that the isomorphism $\mathrm{Lie} \mathbf{H}_g \xrightarrow{\sim} T_{ghg^{-1}}(\mathbf{H}_g)$ was constructed in 2.3(b) using the left multiplication. Therefore the differential form $\omega_{\tilde{\mathbf{G}}_{\mathbf{H}}}(v)$ from 2.3(c) is left-invariant, that is, the restriction of $\omega_{\tilde{\mathbf{G}}_{\mathbf{H}}}(v)$ to each $\mathrm{pr}_{\mathbf{G}/\mathbf{H}}^{-1}([g]) \cong \mathbf{H}_g$ is left invariant.

(b) Instead, we could construct an isomorphism $\mathrm{Lie} \mathbf{H}_g \xrightarrow{\sim} T_{ghg^{-1}}(\mathbf{H}_g)$, using the right multiplication. As a result, for every $v \in V_{\mathbf{G}}$ we would get a right invariant differential form $\omega_{\tilde{\mathbf{G}}_{\mathbf{H}}}^r(v)$ on $\tilde{\mathbf{G}}_{\mathbf{H}}$ such that $\omega_{\tilde{\mathbf{G}}_{\mathbf{H}}}^r(v) = \lambda_{\tilde{\mathbf{G}}_{\mathbf{H}}} \cdot \omega_{\tilde{\mathbf{G}}_{\mathbf{H}}}(v)$ (compare 1.5(b) and 2.2(c)).

(c) Each differential form $\omega_{\tilde{\mathbf{G}}_{\mathbf{H}}}(v)$ is $\mathbf{G}$ -invariant. Indeed, it suffices to show that for every $g \in \mathbf{G}$ and $h \in \mathbf{H}$ the following diagram is commutative

$$\begin{array}{ccc} \mathrm{Lie} \mathbf{H} & \xrightarrow{L_h} & T_h(\mathbf{H}) \\ \mathrm{Ad} g \downarrow & & \downarrow \mathrm{Ad} g \\ \mathrm{Lie} \mathbf{H}_g & \xrightarrow{L_{ghg^{-1}}} & T_{ghg^{-1}}(\mathbf{H}_g). \end{array}$$

But this follows from the identity $g(hx)g^{-1} = (ghg^{-1})(gxg^{-1})$.

Lemma 2.5. *The differential form $a_{\mathbf{H},\mathbf{G}}^*(\omega_{\mathbf{G}}(v))$ on $\tilde{\mathbf{G}}_{\mathbf{H}}$ is equal to $\Delta_{\tilde{\mathbf{G}}_{\mathbf{H}}} \cdot \omega_{\tilde{\mathbf{G}}_{\mathbf{H}}}(v)$, where $\Delta_{\tilde{\mathbf{G}}_{\mathbf{H}}}$ was defined in 1.5(a).*

Proof. Identifying all fibers of $\omega_{\mathbf{G}}(v)$ and $\omega_{\tilde{\mathbf{G}}_{\mathbf{H}}}(v)$ with v as in 2.2(b) and 2.3(b) respectively, we have to show that the Jacobian of the map $a_{\mathbf{H},\mathbf{G}} : \tilde{\mathbf{G}}_{\mathbf{H}} \rightarrow \mathbf{G}$ at $[g, h]$ is $\Delta_{\mathbf{H},\mathbf{G}}(h)$. By the $\mathbf{G}$ -equivariance, we can assume that $g = 1$.

Using 2.3(a), we have two exact sequences of tangent spaces

$$0 \rightarrow \mathrm{Lie} \mathbf{H} \rightarrow T_{[1,h]}(\tilde{\mathbf{G}}_{\mathbf{H}}) \rightarrow \mathrm{Lie} \mathbf{G} / \mathrm{Lie} \mathbf{H} \rightarrow 0$$

$$0 \rightarrow \mathrm{Lie} \mathbf{H} \rightarrow T_h(\mathbf{G}) = \mathrm{Lie} \mathbf{G} \rightarrow \mathrm{Lie} \mathbf{G} / \mathrm{Lie} \mathbf{H} \rightarrow 0.$$

It suffice to show that the differential $da_{\mathbf{H},\mathbf{G}}|_{[1,h]} : T_{[1,h]}(\tilde{\mathbf{G}}_{\mathbf{H}}) \rightarrow T_h(\mathbf{G})$ induces the identity on $\mathrm{Lie} \mathbf{H}$ and the map $\mathrm{Ad} h^{-1} - \mathrm{Id}$ on $\mathrm{Lie} \mathbf{G} / \mathrm{Lie} \mathbf{H}$.

For the first assertion, notice that the restriction of $a_{\mathbf{H},\mathbf{G}}$ to $\mathrm{pr}_{\mathbf{G}/\mathbf{H}}^{-1}([1])$ is the inclusion $\mathbf{H} \hookrightarrow \mathbf{G}$. For the second one, notice that the endomorphism of $\mathrm{Lie} \mathbf{G} / \mathrm{Lie} \mathbf{H}$

induced by $da_{\mathbf{H},\mathbf{G}}|_{[1,h]}$, is induced by the differential of the map $\mathbf{G} \rightarrow \mathbf{G} : g \mapsto ghg^{-1}$ at $g = 1$. Since $\omega_{\mathbf{G}}(v)$ is left invariant, the last differential coincides with the differential of the map $\mathbf{G} \rightarrow \mathbf{G} : g \mapsto (h^{-1}gh)g^{-1}$, which equals $\text{Ad } h^{-1} - \text{Id}$. $\square$

Corollary 2.6. *The open subvariety $\tilde{\mathbf{G}}_{\mathbf{H}}^{\text{reg}} \subset \tilde{\mathbf{G}}_{\mathbf{H}}$ is the largest open subvariety, where the map $a_{\mathbf{H},\mathbf{G}} : \tilde{\mathbf{G}}_{\mathbf{H}} \rightarrow \mathbf{G}$ is étale.*

2.7. The parabolic case. (a) Let $\mathbf{G}$ be connected reductive, $\mathbf{P} \subset \mathbf{G}$ a parabolic subgroup, $\mathbf{M} \subset \mathbf{P}$ a Levi subgroup, and $\mathbf{U} \subset \mathbf{P}$ the unipotent radical.

(b) Let $\mathbf{P}^-$ be the opposite parabolic of $\mathbf{P}$, and let $\mathbf{U}^- \subset \mathbf{P}^-$ be the unipotent radical. Then the multiplication map $m : \mathbf{U}^- \times \mathbf{P} \rightarrow \mathbf{G}$ is an open embedding. Hence the maps $\mathbf{U}^- \rightarrow \mathbf{G}/\mathbf{P} : u \mapsto [u]$ and $j : \mathbf{U}^- \times \mathbf{P} \rightarrow \tilde{\mathbf{G}}_{\mathbf{P}} : (u, x) \mapsto [u, x]$ are open embeddings. In addition, the differential $dm|_{(1,1)} : \text{Lie } \mathbf{U}^- \oplus \text{Lie } \mathbf{P} \rightarrow \text{Lie } \mathbf{G}$ is an isomorphism. Therefore it induces an isomorphism $V_{\mathbf{U}^- \times \mathbf{P}} \xrightarrow{\sim} V_{\mathbf{G}}$.

Claim 2.8. *For every $m \in \mathbf{M}$, we have the identity*

$$\Delta_{\mathbf{P},\mathbf{G}}(m)^2 = (-1)^{\dim \mathbf{U}} \Delta_{\mathbf{M},\mathbf{G}}(m) \cdot \lambda_{\mathbf{P}}(m).$$

Proof. Since both sides of the equality are regular functions on $\mathbf{M}$, we may assume that m is regular semisimple, thus lies in a maximal torus $\mathbf{T} \subset \mathbf{M}$. Now we can decompose $\text{Lie } \mathbf{G}/\text{Lie } \mathbf{M}$ as a sum of $\mathbf{T}$ -eigenspaces. Now the equality follows from the identity $(t-1)^2 = (-1)(t-1)(t^{-1}-1)t$. $\square$

Lemma 2.9. *In the notation of 2.7(b), we have equalities $j^*(\omega_{\tilde{\mathbf{G}}_{\mathbf{P}}}^l(v)) = \omega_{\mathbf{U}^- \times \mathbf{P}}^l(v)$ and $m^*(\omega_{\mathbf{G}}(v)) = \omega_{\mathbf{U}^- \times \mathbf{P}}^r(v)$ for each $v \in V_{\mathbf{G}}$.*

Proof. Since the fiber of each of $j^*(\omega_{\tilde{\mathbf{G}}_{\mathbf{P}}}^l(v))$ and $\omega_{\mathbf{U}^- \times \mathbf{P}}^l(v)$ at 1 is v , for the first equality it suffices to show that $j^*(\omega_{\tilde{\mathbf{G}}_{\mathbf{P}}}^l(v))$ is left $(\mathbf{U}^- \times \mathbf{P})$ -invariant. For the $\mathbf{U}^-$ -invariance, notice that j is $\mathbf{U}^-$ -equivariant, and $\omega_{\tilde{\mathbf{G}}_{\mathbf{P}}}^l(v)$ is $\mathbf{G}$ -invariant (see 2.4(c)). For the $\mathbf{P}$ -invariance notice that $\omega_{\tilde{\mathbf{G}}_{\mathbf{P}}}^l(v)|_{[1] \times \mathbf{P}}$ is left $\mathbf{P}$ -invariant (see 2.4(a)).

For the second equality, one has to show that $m^*(\omega_{\mathbf{G}}(v))$ is right $(\mathbf{U}^- \times \mathbf{P})$ -invariant. Since $\omega_{\mathbf{G}}(v)$ is $\mathbf{G} \times \mathbf{G}$ -invariant, the pullback $m^*(\omega_{\mathbf{G}}(v))$ is right $\mathbf{P}$ -invariant and left $\mathbf{U}^-$ -invariant. Since the group $\mathbf{U}^-$ is unipotent, we have $\lambda_{\mathbf{U}^-} = 1$, thus the differential form $m^*(\omega_{\mathbf{G}}(v))$ is also right $\mathbf{U}^-$ -invariant. $\square$

3. SMOOTH MEASURES, RESTRICTION, AND INDUCTION

From now on, let F be a local non-archimedean field, and let $|\cdot| : F^\times \rightarrow \mathbb{R}^\times$ be the norm map. For every compact analytic subgroup K over F , we denote by δ_K the Haar measure on K with total measure 1.

3.1. Smooth measures. Let X be a smooth analytic variety over F .

(a) Let $C^\infty(X)$ (resp. $C_c^\infty(X)$) be the space of smooth (complex-valued) functions (resp. with compact support), and let $\mathcal{M}(X)$ be the dual space of $C_c^\infty(X)$. Every non-vanishing (F -valued) analytic function f on X induces a smooth function $|f| \in C^\infty(X)$, defined by $|f|(x) := |f(x)|$ for every $x \in X$.

(b) We say that a measure $\chi \in \mathcal{M}(X)$ is *smooth*, and write $\chi \in \mathcal{M}^\infty(X)$, if for every $x \in X$ there exists an open neighborhood $U \subset X$ of x and an analytic isomorphism $\phi : \mathcal{O}_F^n \xrightarrow{\sim} U$ such that $\phi^*(\chi|_U) \in \mathcal{M}(\mathcal{O}_F^n)$ is a multiple of a Haar measure on $\mathcal{O}_F^n$.

(c) By a construction of Weil [We], every non-vanishing top degree differential form ω on X defines a smooth measure $|\omega| \in \mathcal{M}^\infty(X)$.

Namely, for every open analytic embedding $\phi : \mathcal{O}_F^n \xrightarrow{\sim} U \subset X$, the differential form $\phi^*(\omega)$ equals $f dx_1 \wedge \dots \wedge dx_n$ for some non-vanishing analytic function f on $\mathcal{O}_F^n$. Then $|f| \in C_c^\infty(\mathcal{O}_F^n)$, and $|\omega|$ is characterized by the condition that the pullback $\phi^*(|\omega|)$ equals $|f| \delta_{\mathcal{O}_F^n} \in \mathcal{M}^\infty(\mathcal{O}_F^n)$.

Now we give an equivalent (more geometric) definition of smooth measures.

3.2. An alternative description. Let $\mathcal{K}_X^{-1}$ be the line bundle dual to $\mathcal{K}_X$, and let $\Sigma_X \rightarrow X$ be the $F^\times$ -torsor corresponding to $\mathcal{K}_X^{-1}$. Explicitly, Σ_X is a space of pairs (x, a) , where $x \in X$ and a is a non-zero element of the fiber $\mathcal{K}_X^{-1}|_x$ of $\mathcal{K}_X^{-1}$ at x .

Claim 3.3. *The space $\mathcal{M}^\infty(X)$ is canonically identified with the space $C^\infty(\Sigma_X, |\cdot|)$ of smooth functions $f : \Sigma_X \rightarrow \mathbb{C}$ satisfying $f(bx) = |b|f(x)$ for all $b \in F^\times$.*

Proof. Since both $\mathcal{M}^\infty(X)$ and $C^\infty(\Sigma_X, |\cdot|)$ are defined as global sections of certain sheaves on X , we can construct an isomorphism $\mathcal{M}^\infty(X) \xrightarrow{\sim} C^\infty(\Sigma_X, |\cdot|)$ locally on X . Thus we can assume there exists a non-vanishing differential form $\omega \in \Gamma(X, \mathcal{K}_X)$.

Note that the tensor product $s \mapsto s \otimes \omega$ defines an isomorphism of line bundles $\mathcal{K}_X^{-1} \xrightarrow{\sim} \mathcal{O}_X$. Hence it induces an isomorphism $i_\omega : \Sigma_X \xrightarrow{\sim} F^\times \times X$ of the corresponding $F^\times$ -torsors over X . For every $f \in C^\infty(X)$ we denote by $\tilde{f} \in C^\infty(F^\times \times X, |\cdot|)$ the function $(a, x) \mapsto |a|f(x)$. The map $f|\omega| \mapsto i_\omega^*(\tilde{f})$ defines an isomorphism $\mathcal{M}^\infty(X) \xrightarrow{\sim} C^\infty(\Sigma_X, |\cdot|)$, which is independent of ω . $\square$

3.4. Smooth measures with compact support. We denote by $\mathcal{M}_c(X)$ (resp. $\mathcal{H}(X) = \mathcal{M}_c^\infty(X)$) the space of measures (resp. smooth measures) on X with compact support. Moreover, if a group G acts on X , then G acts on the space $\mathcal{H}(X)$, and we denote by $\mathcal{H}(X)^G$ and $\mathcal{H}(X)_G$ the spaces of G -invariants and G -coinvariants, respectively.

3.5. Pullback of smooth measures. (a) Assume that we are given a morphism $f : X \rightarrow Y$ of smooth analytic varieties and an isomorphism $i : f^*(\mathcal{K}_Y) \xrightarrow{\sim} \mathcal{K}_X$ of

line bundles. Then i induces an isomorphism of line bundles $\mathcal{K}_X^{-1} \xrightarrow{\sim} f^*(\mathcal{K}_Y^{-1})$, hence a morphism of $F^\times$ -torsors $(f, i) : \Sigma_X \xrightarrow{\sim} X \times_Y \Sigma_Y \rightarrow \Sigma_Y$.

By Claim 3.3, (f, i) induces a pullback map $(f, i)^* : \mathcal{M}^\infty(Y) \rightarrow \mathcal{M}^\infty(X)$. If, in addition, f is proper, then $(f, i)^*$ induces a pullback $(f, i)^* : \mathcal{H}(Y) \rightarrow \mathcal{H}(X)$.

(b) Note that if $f : X \rightarrow Y$ is a local isomorphism, then the morphism of line bundles $i_f : f^*(\mathcal{K}_Y) \rightarrow \mathcal{K}_X$ from 2.1(b) is an isomorphism. Therefore, by (a), f gives rise to a pullback map $f^* = (f, i_f)^* : \mathcal{M}^\infty(Y) \rightarrow \mathcal{M}^\infty(X)$. Moreover, f induces a pullback map $f^* : \mathcal{H}(Y) \rightarrow \mathcal{H}(X)$, if in addition f is proper.

In particular, if $X \hookrightarrow Y$ is an open (resp. open and closed) embedding, then we have a restriction map $\text{res} : \mathcal{M}^\infty(Y) \rightarrow \mathcal{M}^\infty(X)$ (resp. $\text{res} : \mathcal{H}(Y) \rightarrow \mathcal{H}(X)$).

The following simple lemma is basic for what follows.

Lemma 3.6. (a) *Let $f : X \rightarrow Y$ be a smooth map between smooth analytic varieties. Then the pushforward map $f_! : \mathcal{M}_c(X) \rightarrow \mathcal{M}_c(Y)$ satisfies $f_!(\mathcal{H}(X)) \subset \mathcal{H}(Y)$, that is, $f_!(h) \in \mathcal{H}(Y)$ for every $h \in \mathcal{H}(X)$.*

(b) *Let G be an analytic group, and let $f : X \rightarrow Y$ be a principal G -bundle. Then the map $f_!$ induces an isomorphism $\mathcal{H}(X)_G \xrightarrow{\sim} \mathcal{H}(Y)$.*

Proof. (a) The question is local in X and Y , so we may assume that $X = \mathcal{O}_F^{n+m}$, $Y = \mathcal{O}_F^n$, f is the projection, and $h = \delta_X$. In this case, $f_!(h) = \delta_Y$.

(b) The assertion is local in Y , and f is locally trivial, so we may assume that $X = G \times Y$. In this case, for every compact open subgroup $K \subset G$, the map $\mathcal{H}(Y) \rightarrow \mathcal{H}(X)_G$ defined by $h \mapsto [h \boxtimes \delta_K]$ is the inverse of $f_! : \mathcal{H}(X)_G \xrightarrow{\sim} \mathcal{H}(Y)$. $\square$

3.7. The induced space. (a) Let G an analytic group, $H \subset G$ a closed analytic subgroup, and let H act on a smooth analytic variety X . Then the product $G \times X$ is equipped with an action of $G \times H$ defined by $(g, h)(g', x) := (gg'h^{-1}, h(x))$, and the quotient $\text{Ind}_H^G(X) := G \times^H X$ is smooth and equipped with an action of G .

(b) Consider the diagram $\text{Ind}_H^G(X) \xleftarrow{p_1} G \times X \xrightarrow{p_2} X$, where p_1 and p_2 are the natural projections. Since p_1 is a G -equivariant H -bundle, while p_2 is an H -equivariant G -bundle, Lemma 3.6(b) implies that we have a natural isomorphism

$$\varphi_H^G := (p_2)_! \circ (p_1)_!^{-1} : \mathcal{H}(\text{Ind}_H^G(X))_G \xleftarrow{\sim} \mathcal{H}(G \times X)_{G \times H} \xrightarrow{\sim} \mathcal{H}(X)_H.$$

(c) By construction, the isomorphism φ_H^G from (b) can be characterized as the unique map $\varphi_H^G : \mathcal{H}(\text{Ind}_H^G(X))_G \rightarrow \mathcal{H}(X)_H$ such that the composition $\varphi_H^G \circ (p_1)_! : \mathcal{H}(G \times X)_{G \times H} \rightarrow \mathcal{H}(X)_H$ is equal to $(p_2)_!$.

(d) Assume that H acts on X trivially. Then $\text{Ind}_H^G(X) = (G/H) \times X$, and the natural projection $\text{pr} : (G/H) \times X \rightarrow X$ satisfies $\text{pr} \circ p_1 = p_2$. Thus, by (c), the map φ_H^G obtained from $\text{pr}_! : \mathcal{H}((G/H) \times X) \rightarrow \mathcal{H}(X)$.

(e) Assume that H is a retract of G , and let $p : G \rightarrow H$ be a homomorphism such that $p|_H = \text{Id}_H$. Then p induces a map $p : \text{Ind}_H^G(X) \rightarrow \text{Ind}_H^H(X) = X$ such that $p \circ p_1 = p_2$. Thus, by (c), the map φ_H^G is obtained from $p_! : \mathcal{H}(\text{Ind}_H^G(X)) \rightarrow \mathcal{H}(X)$.

3.8. Notation. For every (smooth) algebraic variety $\mathbf{X}$ over F we denote by X the corresponding (smooth) analytic variety $\mathbf{X}(F)$. In particular, we have $G = \mathbf{G}(F)$, $\tilde{G}_H = \tilde{\mathbf{G}}_H(F)$, etc. We also denote by $a_{H,G} : \tilde{G}_H \rightarrow G$, $\Delta_{H,G} : H \rightarrow F$, $\lambda_G : G \rightarrow F^\times$, etc. the maps induced by $a_{\mathbf{H},\mathbf{G}}$, $\Delta_{\mathbf{H},\mathbf{G}}$ and $\lambda_{\mathbf{G}}$, respectively.

3.9. The restriction map. In the situation of 2.7, set $\tilde{G} := \tilde{G}_P$ and $a := a_{P,G}$.

(a) In 2.3(d) we constructed an $\text{Ad } G$ -equivariant isomorphism $\tilde{i} : a^*(\mathcal{K}_G) \xrightarrow{\sim} \mathcal{K}_{\tilde{G}}$, which by 3.5(a) defines a pullback map $\tilde{a}^* = (a, \tilde{i})^* : \mathcal{M}^\infty(G) \rightarrow \mathcal{M}^\infty(\tilde{G})$.

Explicitly, for every $v \in V_G$ and $f \in C^\infty(G)$, the map $\tilde{a}^*$ is given by the formula $\tilde{a}^*(f|\omega_G(v)|) := a^*(f)|\omega_{\tilde{G}}^l(v)|$. Moreover, since $a : \tilde{G} \rightarrow G$ is proper, the map $\tilde{a}^*$ induces a G -equivariant map $\tilde{a}^* : \mathcal{H}(G) \rightarrow \mathcal{H}(\tilde{G})$.

(b) Since $(\mathbf{G}/\mathbf{P})(F)$ equals $\mathbf{G}(F)/\mathbf{P}(F) = G/P$ (see [Bo, Prop 20.5]), the set $\tilde{G} = \tilde{\mathbf{G}}(F)$ equals $\text{Ind}_P^G(P)$. Hence, by 3.7(b), we have a natural isomorphism

$$\varphi_P^G : \mathcal{H}(\tilde{G})_G = \mathcal{H}(\text{Ind}_P^G(P))_G \xrightarrow{\sim} \mathcal{H}(P)_P.$$

We denote by $\text{Res}_P^G : \mathcal{H}(G)_G \rightarrow \mathcal{H}(P)_P$ the composition map $\varphi_P^G \circ \tilde{a}^*$.

(c) Let $p : \mathbf{P} \rightarrow \mathbf{M}$ be the projection (see Lemma 1.9). Since p is smooth, the induced map $p : P \rightarrow M$ is smooth. Thus, p induces a map $p_! : \mathcal{H}(P)_P \rightarrow \mathcal{H}(M)_M$ (see Lemma 3.6(a)), and we denote by $\text{Res}_{P,M}^G : \mathcal{H}(G)_G \rightarrow \mathcal{H}(M)_M$ the composition $p_! \circ \text{Res}_P^G$.

3.10. Generalized functions and induction. (a) Let X be a smooth analytic variety over F , and let $\widehat{C}(X) := \text{Hom}_{\mathbb{C}}(\mathcal{H}(X), \mathbb{C})$ be the space generalized functions on X . Then the space $\widehat{C}(X)$ is equipped with a *weak topology*, which is the coarsest topology such that for every $h \in \mathcal{H}(X)$ and $a \in \mathbb{C}$ the set $\{\lambda \in \widehat{C}(X) \mid \lambda(h) \neq a\}$ is open.

(b) For a group G acting on X , we denote by $\widehat{C}^G(X) \subset \widehat{C}(X)$ the subspace of invariant generalized functions with the induced topology. Equivalently, $\widehat{C}^G(X)$ is the linear dual space of $\mathcal{H}(X)_G$ equipped with the weak topology.

(c) By Lemma 3.6(a), every smooth map $f : X \rightarrow Y$ induces a continuous map $f^* : \widehat{C}(Y) \rightarrow \widehat{C}(X)$ dual to $f_!$. If, in addition, f is G -equivariant, then f^* induces a continuous map $\widehat{C}^G(Y) \rightarrow \widehat{C}^G(X)$.

(d) In the situation of 3.7, we have a linear homeomorphism $\widehat{C}^H(X) \xrightarrow{\sim} \widehat{C}^H(\text{Ind}_H^G(X))$, dual to φ_H^G .

(e) In the situation of 3.9, we denote by $\text{Ind}_{P;M}^G : \widehat{C}^M(M) \rightarrow \widehat{C}^G(G)$ (resp. $\text{Ind}_P^G : \widehat{C}^P(P) \rightarrow \widehat{C}^G(G)$) the map dual to $\text{Res}_{P;M}^G$ (resp. Res_P^G).

(f) Let $\mathbf{H}$ be an algebraic group over F . Then to every admissible representation π of $H = \mathbf{H}(F)$ we can associate its character $\chi_\pi \in \widehat{C}^H(H)$.

4. RELATION TO CHARACTERS OF INDUCED REPRESENTATIONS

Assume that we are in the situation of 3.9.

Proposition 4.1. *Let τ be an admissible representation of P , and let $\pi = \text{Ind}_P^G(\tau)$ be the induced representation. Then we have the equality $\chi_\pi = \text{Ind}_P^G(\chi_\tau)$.*

To prove the result, we will compute both χ_π and $\text{Ind}_P^G(\chi_\tau)$ explicitly.

4.2. Notation. (a) Let $K \subset G$ be a compact open subgroup, set $K_P := K \cap P$, and let μ^K (resp. μ^{K_P}) be the left invariant Haar measure on G (resp. P) normalized by the condition that $\mu_K(K) = 1$ (resp. $\mu^{K_P}(K_P) = 1$).

(b) Set $\tilde{K} := K \overset{K_P, \text{Ad}}{\times} P = \text{Ind}_{K_P}^K(P)$. Then $\tilde{K} \subset \tilde{G}$ is an open and closed subset, and we set

$$\text{Res}_{K_P}^K : \mathcal{H}(G)_K \xrightarrow{\tilde{a}^*} \mathcal{H}(\tilde{G})_K \xrightarrow{\text{res}} \mathcal{H}(\tilde{K})_K \xrightarrow{\varphi_{K_P}^K} \mathcal{H}(P)_{K_P},$$

where $\tilde{a}^*$ was defined in 3.9(a) and $\varphi_{K_P}^K$ was defined in 3.7(b).

(c) For every $h \in \mathcal{H}(G)^K$, we define $f := h/\mu^K \in C_c^\infty(G)$, $f_P := f|_P \in C_c^\infty(P)$ and $h_P := f_P \mu^{K_P} \in \mathcal{H}(P)$.

(d) For every $g \in G$, we set $K_g := gKg^{-1}$, $h_g := (\text{Ad } g^{-1})^*(h) \in \mathcal{H}(G)^{K_g}$, $K_{g,P} := K_g \cap P$ and $h_{g,P} := (h_g)_P$ (see (c)).

(e) Fix a set of representatives $A \subset G$ of double classes $P \backslash G / K$. The set A is finite, because $P \backslash G$ is compact.

Lemma 4.3. *In the notation of 4.2, we have equalities:*

- (a) $\text{Res}_P^G([h]) = \sum_{g \in A} \text{Res}_{K_{g,P}}^{K_g}([h_g])$ in $\mathcal{H}(P)_P$;
- (b) $\text{Res}_{K_P}^K([h]) = [h_P]$ in $\mathcal{H}(P)_{K_P}$;
- (c) $\text{Res}_P^G([h]) = \sum_{g \in A} [h_{g,P}]$ in $\mathcal{H}(P)_P$.

Proof. (a) By definition, $\text{Res}_P^G([h]) = \varphi_P^G([\tilde{a}^*(h)])$. Notice that the decomposition $G = \sqcup_{g \in A} Kg^{-1}P$ into open and closed subsets induces a decomposition $\tilde{G} = \sqcup_g \tilde{G}_g$, where $\tilde{G}_g := Kg^{-1}P \overset{P}{\times} P$. Therefore we get a decomposition $\tilde{a}^*(h) = \sum_g \{h\}_g$, where $\{h\}_g := \tilde{a}^*(h)|_{\tilde{G}_g} \in \mathcal{H}(\tilde{G}_g) \subset \mathcal{H}(\tilde{G})$. Thus it remains to show that for every $g \in G$, we have the equality $\varphi_P^G([\{h\}_g]) = \text{Res}_{K_{g,P}}^{K_g}([h_g])$.

Note that $g\widetilde{G}_g = gKg^{-1}P \overset{P}{\times} P = K_g \overset{K_{g,P}}{\times} P = \widetilde{K}_g$ and $(g^{-1})^*(\{h\}_g) = \widetilde{a}^*(h_g)|_{\widetilde{K}_g}$. Using the identity $[\{h\}_g] = [(g^{-1})^*(\{h\}_g)] \in \mathcal{H}(\widetilde{G})_G$, the equality $\varphi_P^G([\{h\}_g]) = \text{Res}_{K_{g,P}}^{K_g}([h_g])$ can be rewritten as $\varphi_P^G([\widetilde{a}^*(h_g)|_{\widetilde{K}_g}]) = \text{Res}_{K_{g,P}}^{K_g}([h_g])$.

The latter equality follows from the fact that the diagram

$$\begin{array}{ccc} \mathcal{H}(\widetilde{G}) & \xrightarrow{\varphi_P^G} & \mathcal{H}(P)_P \\ (1) \uparrow & & (2) \uparrow \\ \mathcal{H}(\widetilde{K}_g) & \xrightarrow{\varphi_{K_{g,P}}^{K_g}} & \mathcal{H}(P)_{K_{g,P}}, \end{array}$$

where (1) is the natural inclusion, while (2) is the natural projection, is commutative.

(b) Recall that $\varphi_{K_P}^K$ is a composition $\mathcal{H}(\widetilde{K})_K \xleftarrow{\sim} \mathcal{H}(K \times P)_{K \times K_P} \xrightarrow{\sim} \mathcal{H}(P)_{K_P}$, corresponding to the diagram $\widetilde{K} \xleftarrow{p_1} K \times P \xrightarrow{p_2} P$. Since element $\delta_K \boxtimes h_P$ of $\mathcal{H}(K \times P)$ satisfies $(p_2)_!(\delta_K \boxtimes h_P) = h_P$, it suffices to show the equality $(p_1)_!(\delta_K \boxtimes h_P) = \widetilde{a}^*(h)|_{\widetilde{K}}$.

Note that $\widetilde{a}^*(h) = a^*(f)\widetilde{a}^*(\mu^K)$, while $\delta_K \boxtimes h_P = (1_K \boxtimes f_P)(\delta_K \times \mu^{K_P})$. Moreover, since f is $\text{Ad } K$ -invariant, we have $p_1^*(a^*(f)) = 1_K \boxtimes f_P$. Thus it remains to show the equality $(p_1)_!(\delta_K \boxtimes \mu^{K_P}) = \widetilde{a}^*(\mu^K)|_{\widetilde{K}}$.

Fix $0 \neq v \in V_G$ and $0 \neq v' \in V_P$. Using the identities $|\omega_G(v)| = |\omega_G(v)|(K) \cdot \mu^K$ and $|\omega_P^l(v')| = |\omega_P^l(v')|(K_P) \cdot \mu^{K_P}$, it remains to show the equality

$$(4.1) \quad (p_1)_!(|\omega_K(v)| \boxtimes |\omega_P^l(v')|) = |\omega_P^l(v')|(K_P) \cdot |\omega_{\widetilde{K}}(v)|,$$

where we set $\omega_{\widetilde{K}}(v) = \omega_{\widetilde{G}}(v)|_{\widetilde{G}}$. Using the notation of 2.7(b), we set $K^- := K \cap U^-$, and consider the open embeddings $j : K^- \times P \hookrightarrow \widetilde{K}$, $m : K^- \times K_P \hookrightarrow K$ and $K^- \times P \hookrightarrow G$. Since p_1 is K -equivariant, it remains to show the restriction of the equality (4.1) to $K^- \times P$ under j .

Note that $p_1^{-1}(K^- \times P) = K^- \times K_P \times P$, while $j^*(\omega_{\widetilde{K}}(v)) = \omega_{K^- \times P}^l(v)$ and $m^*(\omega_K(v)) = \omega_{K^- \times K_P}^r(v)$ (by Lemma 2.9). Thus, the assertion follows from the fact that $|\omega_{K^- \times K_P}^r(v)| = |\omega_{K^- \times K_P}^l(v)|$, because $K^- \times K_P$ is compact.

(c) follows immediately from (a) and (b). $\square$

4.4. Proof of Proposition 4.1. We have to show that for every $h \in \mathcal{H}(G)$ we have $\chi_\pi(h) = \chi_\tau(\text{Res}_P^G(h))$. Choose an open compact subgroup $K \subset G$ such that h is $K \times K$ -invariant. Then $h \in \mathcal{H}(G)^K$, so by Lemma 4.3 we have to show that

$$(4.2) \quad \text{Tr}(h, \pi) = \sum_{g \in A} \text{Tr}(h_{g,P}, \tau).$$

Though the result is well-known and is an immediate generalization of the corresponding result for finite groups, we sketch the argument for completeness.

Let W (resp. V) be the space of τ (resp. π). Then V^K is the space of functions $f : G \rightarrow W$ satisfying $f(xyk) = \tau(x)(f(y))$ for all $x \in P$, $y \in G$ and $k \in K$. For every $g \in G$ we denote by $V_g^K \subset V^K$ the subspace of functions $f : G \rightarrow W$ from V^K , supported on PgK . Then $V^K = \bigoplus_{g \in A} V_g^K$.

For every $g \in G$, consider the endomorphism $h_{\{g\}} : V_g^K \hookrightarrow V^K \xrightarrow{h} V^K \twoheadrightarrow V_g^K$, induced by h . Then $\text{Tr}(h, \pi) = \sum_{g \in A} \text{Tr}(h_{\{g\}})$, so it suffices to show the equality $\text{Tr}(h_{\{g\}}) = \text{Tr}(h_{g,P}, W)$ for every $g \in G$.

Since $gK = K_g g$, the map $f \mapsto f(g)$ induces a linear isomorphism $V_g^K \xrightarrow{\sim} W^{K_{g,P}}$. It remains to show that this isomorphism identifies $h_{\{g\}} \in \text{End } V_g^K$ with $\tau(h_{g,P}) \in \text{End } W^{K_{g,P}}$, that is, for every $f \in V_g^K$ we have the equality $(h(f))(g) = \tau(h_{g,P})(f(g))$.

We claim that the latter equality holds for every right K -invariant $h \in \mathcal{H}(G)$. Indeed, we may assume that $h = \delta_{xK}$ for some $x \in G$. Set $x_g := gxg^{-1}$. Then $h_g = \delta_{x_g K_g}$, and $(h(f))(g) = f(gx) = f(x_g g)$. Assume first that $x_g \in PK_g$. Then $x_g K_g = x' K_g$ for some $x' \in P$. In this case, we have $h_{g,P} = \delta_{x' K_{g,P}}$, and

$$h(f)(g) = f(x_g g) = f(x' g) = \tau(x')(f(g)) = \tau(h_{g,P})(f(g)).$$

Finally, if $x_g \notin PK_g$, then $h(f)(g) = f(x_g g) = 0$, and $h_{g,P} = 0$. $\square$

4.5. Parabolic induction. Let ρ be an admissible representation of M . Recall that a non-normalized parabolic induction $\pi = \text{Ind}_{P,M}^G(\rho)$ is the induced representation $\text{Ind}_P^G(\tau)$, where $\tau \in \text{Rep}(P)$ is the inflation of ρ .

Corollary 4.6. *We have the equality of characters $\chi_\pi = \text{Ind}_{P,M}^G(\chi_\rho)$.*

Proof. Since the character of the inflation $\chi_\tau \in \widehat{C}^P(P)$ equals $p^*(\chi_\rho)$, the assertion follows from Proposition 4.1. $\square$

5. RESTRICTION TO AN EQUAL RANK SUBGROUP

5.1. Smooth measures with relatively compact support.

(a) Let $f : X \rightarrow Z$ be a morphism of analytic varieties over F , where X is smooth. We denote by $\mathcal{H}(X/Z) \subset \mathcal{M}^\infty(X)$ the subspace consisting of measures, whose support is proper over Z . In particular, $\mathcal{H}(X/X) = \mathcal{M}^\infty(X)$. Notice that $\mathcal{H}(X) \subset \mathcal{H}(X/Z)$, if Z is Hausdorff, and $\mathcal{H}(X) = \mathcal{H}(X/Z)$, if Z is compact.

(b) Every smooth morphism $f : X \rightarrow Y$ over Z induces a canonical push-forward map $f_! : \mathcal{H}(X/Z) \rightarrow \mathcal{H}(Y/Z)$. Indeed, we can construct the map $f_!$ locally on Z , thus may assume that Z is compact. Now the assertion follows from Lemma 3.6.

(c) Assume that we are in the situation of 3.7, and that H acts on X over Z , that is, the map $f : X \rightarrow Z$ is H -equivariant with respect to the trivial action of H on Z . Then the arguments of (b) and 3.7 imply that we have a natural isomorphism $\varphi_H^G : \mathcal{H}(\text{Ind}_H^G(X)/Z)_G \xrightarrow{\sim} \mathcal{H}(X/Z)_H$.

(d) Assume that we are given a commutative diagram of analytic spaces over F

$$\begin{array}{ccc} X & \longrightarrow & Y \\ b \downarrow & & \downarrow \\ X' & \longrightarrow & Y', \end{array}$$

such that X and X' are smooth, b is a local isomorphism, and the induced map $X \rightarrow X' \times_{Y'} Y$ is proper. Then the pullback $b^* : \mathcal{M}^\infty(X') \rightarrow \mathcal{M}^\infty(X)$ (see 3.5(b)) satisfies $b^*(\mathcal{H}(X'/Y')) \subset \mathcal{H}(X/Y)$.

(e) An important particular case of (d) is when the diagram is Cartesian and $Y \rightarrow Y'$ is an open embedding. In this case, $b : X \rightarrow X'$ is an open embedding as well, and we denote b^* by res and call it *the restriction map*.

5.2. The algebraic case. (a) Let $\mathbf{H} \subset \mathbf{G}$ be a closed subgroup. Since the projection map $\mathbf{G} \rightarrow \mathbf{G}/\mathbf{H}$ is smooth, every G -orbit in $(\mathbf{G}/\mathbf{H})(F)$ is open. Therefore $G/H = \mathbf{G}(F)/\mathbf{H}(F)$ is an open and closed subset of $(\mathbf{G}/\mathbf{H})(F)$.

(b) Let $\mathbf{H}$ act on a smooth algebraic variety $\mathbf{X}$. Then, by (a), the induced space $\text{Ind}_H^G(X)$ is an open and closed subset of $(\text{Ind}_{\mathbf{H}}^{\mathbf{G}}(\mathbf{X}))(F)$. Hence we have a natural restriction map $\text{res} : \mathcal{H}((\text{Ind}_{\mathbf{H}}^{\mathbf{G}}(\mathbf{X}))(F))_G \rightarrow \mathcal{H}(\text{Ind}_H^G(X))_G$, and we denote the composition $\varphi_H^G \circ \text{res} : \mathcal{H}((\text{Ind}_{\mathbf{H}}^{\mathbf{G}}(\mathbf{X}))(F))_G \rightarrow \mathcal{H}(X)_H$ simply by φ_H^G .

(c) Assume that $\mathbf{H}$ acts on $\mathbf{X}$ over $\mathbf{Z}$ (compare 5.1(c)). Then, generalizing (b), we have a map $\varphi_H^G : \mathcal{H}((\text{Ind}_{\mathbf{H}}^{\mathbf{G}}(\mathbf{X}))(F)/Z)_G \rightarrow \mathcal{H}(X/Z)_H$.

5.3. Notation. Let $\mathbf{K} \subset \mathbf{H}$ be two closed subgroups of $\mathbf{G}$.

(a) We set $\mathcal{H}(H)^{\text{reg}/G} := \mathcal{H}(H^{\text{reg}/G}/c_H^{\text{reg}/G})$, where $H^{\text{reg}/G} = \mathbf{H}^{\text{reg}/\mathbf{G}}(F)$ and $c_H^{\text{reg}/G} = \mathbf{c}_{\mathbf{H}}^{\text{reg}/\mathbf{G}}(F)$ (see 1.2(b)). Similarly, we set $\mathcal{H}(\tilde{G}_H)^{\text{reg}} := \mathcal{H}(\tilde{G}_H^{\text{reg}}/c_H^{\text{reg}/G})$ and $\mathcal{H}(\tilde{H}_K)^{\text{reg}/G} := \mathcal{H}(\tilde{H}_K^{\text{reg}/G}/c_K^{\text{reg}/G})$ (see 1.5(e)).

(b) By a combination of 5.1(a) and 5.1(e), we have a restriction map

$$\text{res} : \mathcal{H}(H) \hookrightarrow \mathcal{H}(H/c_H) \rightarrow \mathcal{H}(H)^{\text{reg}/G}.$$

(c) Recall that the map $a_{\mathbf{H},\mathbf{G}} : \tilde{\mathbf{G}}_{\mathbf{H}}^{\text{reg}} \rightarrow \mathbf{G}$ is $\mathbf{G}$ -equivariant and étale (see Corollary 2.6), thus the corresponding map $a_{H,G} : \tilde{G}_H^{\text{reg}} \rightarrow G$ is a local isomorphism. Thus we have a pullback map $a_{H,G}^* : \mathcal{M}^\infty(G) \rightarrow \mathcal{M}^\infty(\tilde{G}_H^{\text{reg}})$ (see 3.5(b)).

5.4. The restriction map. Assume that we are in the situation of 1.6.

(a) Since the morphism of algebraic varieties $\iota_{\mathbf{H}, \mathbf{G}} : \tilde{\mathbf{G}}_{\mathbf{H}}^{\text{reg}} \rightarrow \mathbf{G} \times_{\mathbf{c}_{\mathbf{G}}} \mathbf{c}_{\mathbf{H}}^{\text{reg}/\mathbf{G}}$ is finite (by Corollary 1.11), the corresponding morphism of analytic varieties $\iota_{H, G} : \tilde{G}_H^{\text{reg}} \rightarrow G \times_{c_G} c_H^{\text{reg}/G}$ is proper. Thus, by 5.3(c) and 5.1(d) we have a pullback map $a_{H, G}^* : \mathcal{H}(G)_G \rightarrow \mathcal{H}(G/c_G)_G \rightarrow \mathcal{H}(\tilde{G}_H)_G^{\text{reg}}$.

(b) We denote by $R_H^G : \mathcal{H}(G)_G \rightarrow \mathcal{H}(\tilde{G}_H)_G^{\text{reg}} \rightarrow \mathcal{H}(H)_H^{\text{reg}/G}$ the composition of the map $a_{H, G}^*$ from (a) and the map φ_H^G from 5.2(c).

5.5. Set up. Assume that we are in the situation of 3.9, and that $\mathbf{H}$ is a connected equal rank subgroup of $\mathbf{M}$, hence also of $\mathbf{G}$.

(a) Then $\mathbf{c}_{\mathbf{H}}^{\text{reg}/\mathbf{G}} \subset \mathbf{c}_{\mathbf{H}}^{\text{reg}/\mathbf{M}}$ is an open subscheme (see 1.2(c)), thus we have defined a restriction map $\text{res} : \mathcal{H}(H)_H^{\text{reg}/M} \rightarrow \mathcal{H}(H)_H^{\text{reg}/G}$ (see 5.1(e)).

(b) Using 1.2(c) again, we conclude that $\mathbf{H}^{\text{reg}/\mathbf{G}} \subset \mathbf{P}^{\text{reg}/\mathbf{G}}$. Thus the restriction of $\Delta_{\mathbf{P}, \mathbf{G}}$ to $\mathbf{H}^{\text{reg}/\mathbf{G}}$ is non-vanishing. Hence $\Delta_{\mathbf{P}, \mathbf{G}}$ gives rise to a smooth function $|\Delta_{P, G}| \in \mathcal{M}^\infty(H^{\text{reg}/G})$.

Lemma 5.6. *In the situation of 5.5, the following diagram is commutative*

$$(5.1) \quad \begin{array}{ccc} \mathcal{H}(G)_G & \xrightarrow{R_H^G} & \mathcal{H}(H)_H^{\text{reg}/G} \\ \text{Res}_{P, M}^G \downarrow & & \uparrow |\Delta_{P, G}| \cdot \text{res} \\ \mathcal{H}(M)_M & \xrightarrow{R_H^M} & \mathcal{H}(H)_H^{\text{reg}/M} \end{array}$$

Proof. The assertion follows from a rather straightforward diagram chase. Namely, using the inclusion $H^{\text{reg}/G} \subset M^{\text{reg}/G}$, we observe that diagram (5.1) decomposes as

$$(5.2) \quad \begin{array}{ccccc} \mathcal{H}(G)_G & \xrightarrow{R_M^G} & \mathcal{H}(M)_M^{\text{reg}/G} & \xrightarrow{R_H^M} & \mathcal{H}(H)_H^{\text{reg}/G} \\ \text{Res}_{P, M}^G \downarrow & & \uparrow |\Delta_{P, G}| \cdot \text{res} & & \uparrow |\Delta_{P, G}| \cdot \text{res} \\ \mathcal{H}(M)_M & \xlongequal{\quad} & \mathcal{H}(M)_M & \xrightarrow{R_H^M} & \mathcal{H}(H)_H^{\text{reg}/M} \end{array}$$

Since the right inner square of (5.2) is commutative by functoriality, it remains to show the commutativity of the left inner square.

Observe that the diagram

$$(5.3) \quad \begin{array}{ccccc} \mathcal{H}(P)_P & \xrightarrow{\text{res}} & \mathcal{H}(P)_P^{\text{reg}/G} & \xrightarrow{a_{M, P}^*} & \mathcal{H}(\tilde{P}_M)_P^{\text{reg}/G} \\ p! \downarrow & & p! \downarrow & & \varphi_M^P \downarrow \\ \mathcal{H}(M)_M & \xrightarrow{\text{res}} & \mathcal{H}(M)_M^{\text{reg}/G} & \xlongequal{\quad} & \mathcal{H}(M)_M^{\text{reg}/G} \end{array}$$

is commutative. Indeed, the left inner square of (5.3) is commutative by Lemma 1.9(d), and the right inner square is commutative, because $\varphi_M^P = p_!$ (see 3.7(e)) and $a_{P,M}^*$ is an isomorphism (by Lemma 1.9(e)).

Therefore the left inner square of (5.2) decomposes as

$$(5.4) \quad \begin{array}{ccccc} \mathcal{H}(G)_G & \xlongequal{\quad} & \mathcal{H}(G)_G & \xlongequal{\quad} & \mathcal{H}(G)_G \\ \tilde{a}_{P,G}^* \downarrow & & a_{M,G}^* \downarrow & & R_M^G \downarrow \\ \mathcal{H}(\tilde{G}_P)_G & \xrightarrow{|\Delta_{P,G}| \cdot a_{M,P}^*} & \mathcal{H}(\tilde{G}_M)_G^{\text{reg}} & \xrightarrow{\varphi_M^G} & \mathcal{H}(M)_M^{\text{reg}/G} \\ \varphi_P^G \downarrow & & \varphi_P^G \downarrow & & \parallel \\ \mathcal{H}(P)_P & \xrightarrow{|\Delta_{P,G}| \cdot a_{M,P}^*} & \mathcal{H}(\tilde{P}_M)_G^{\text{reg}/G} & \xrightarrow{\varphi_M^P} & \mathcal{H}(M)_M^{\text{reg}/G}. \end{array}$$

We claim that all inner squares of (5.4) are commutative. Indeed, the top right square is commutative by the definition of R_M^G , the bottom left square is commutative by the functoriality of φ_P^G , and the bottom right square is commutative by the equality $\varphi_M^P \circ \varphi_P^G = \varphi_M^G$.

Finally, the commutativity of the top left square of (5.4) follows from the equality $a_{M,G} = a_{M,P} \circ a_{P,G}$ (see 1.5(d)) and Lemma 2.5. $\square$

6. NORMALIZED INDUCTION, INDEPENDENCE OF $\mathbf{P}$, AND STABILITY

6.1. Normalized restriction and induction.

(a) Recall that in the construction of the restriction map $\text{Res}_{P,M}^G$ in 3.9(c) we used the isomorphism $\mathcal{O}_{\tilde{\mathbf{G}}} \otimes_F V_{\mathbf{G}} \xrightarrow{\sim} \mathcal{K}_{\tilde{\mathbf{G}}} : v \mapsto \omega_{\tilde{\mathbf{G}}}(v)$ (see 2.3(c)). Instead we could use the isomorphism $\mathcal{O}_{\tilde{\mathbf{G}}} \otimes_F V_{\mathbf{G}} \xrightarrow{\sim} \mathcal{K}_{\tilde{\mathbf{G}}}^r : v \mapsto \omega_{\tilde{\mathbf{G}}}^r(v)$ (see 2.4(b)). Since $\omega_{\tilde{\mathbf{G}}}^r(v) = \lambda_{\tilde{\mathbf{G}}} \cdot \omega_{\tilde{\mathbf{G}}}(v)$, the resulting restriction map would be $|\lambda_P| \cdot \text{Res}_{P,M}^G$.

(b) We denote by $r_{P,M}^G : \mathcal{H}(G)_G \rightarrow \mathcal{H}(M)_M$ the map $|\lambda_P|^{1/2} \cdot \text{Res}_{P,M}^G$, and call it the *normalized restriction map*. Let $i_{P,M}^G : \hat{C}^M(M) \rightarrow \hat{C}^G(G)$ be the dual map of $r_{P,M}^G$, called the *normalized induction map*. Explicitly, $i_{P,M}^G(\chi) = \text{Ind}_{P,M}^G(|\lambda_P|^{1/2} \cdot \chi)$ for every $\chi \in \hat{C}^M(M)$.

(c) For an admissible representation ρ of M , we denote by $\underline{i}_{P,M}^G(\rho)$ the representation $\underline{\text{Ind}}_{P,M}^G(\rho \otimes |\lambda_P|^{1/2})$ of G and call it the *normalized parabolic induction*.

6.2. Remark. If $\mathbf{G}$ is semisimple and simply connected, and $\mathbf{P}$ is a Borel subgroup of $\mathbf{G}$, then the normalized restriction map $r_{P,M}^G$ has a geometric interpretation.

Indeed, in this case, the homomorphism $\lambda_{\mathbf{P}} : \mathbf{P} \rightarrow \mathbb{G}_m$ has a unique square root $\lambda_{\mathbf{P}}^{1/2} : \mathbf{P} \rightarrow \mathbb{G}_m$. Furthermore, $\lambda_{\mathbf{P}}^{1/2}$ gives rise to a morphism $\lambda_{\tilde{\mathbf{G}}}^{1/2} : \tilde{\mathbf{G}} \rightarrow \mathbb{G}_m$ (by

1.5(a), hence to an isomorphism $\mathcal{O}_{\tilde{\mathbf{G}}} \otimes_F V_{\mathbf{G}} \xrightarrow{\sim} \mathcal{K}_{\tilde{\mathbf{G}}} : v \mapsto \lambda_{\tilde{\mathbf{G}}}^{1/2} \cdot \omega_{\tilde{\mathbf{G}}}(v)$, and $r_{P;M}^G$ is obtained from this isomorphism by the construction 3.9.

The following result follows immediately from Corollary 4.6.

Corollary 6.3. *We have the equality of characters $\chi_{i_{P;M}^G(\rho)} = i_{P;M}^G(\chi_\rho)$.*

Next, we write a version of Lemma 5.6 for the normalized restriction.

Corollary 6.4. *In the situation of 5.5, the following diagram is commutative*

$$(6.1) \quad \begin{array}{ccc} \mathcal{H}(G)_G & \xrightarrow{R_H^G} & \mathcal{H}(H)_H^{\text{reg}/G} \\ r_{P;M}^G \downarrow & & \uparrow |\Delta_{M,G}|^{1/2} \cdot \text{res} \\ \mathcal{H}(M)_M & \xrightarrow{R_H^M} & \mathcal{H}(H)_H^{\text{reg}/M}. \end{array}$$

Proof. By Claim 2.8, we have the identity $|\Delta_{M,G}|^{1/2} = |\Delta_{P,G}| \cdot |\lambda_P|^{-1/2}$, so the assertion follows from Lemma 5.6. $\square$

Lemma 6.5. *The restriction map $\text{res} : \mathcal{H}(M)_M \rightarrow \mathcal{H}(M)_M^{\text{reg}/G}$ from 5.3(b) is injective.*

Proof. By definition, the connected center $\mathbf{Z}(\mathbf{M})^0 \subset \mathbf{Z}(\mathbf{M})$ acts on $\text{Lie } \mathbf{G} / \text{Lie } \mathbf{M}$ by a direct sum of non-trivial characters. Therefore for every $m \in \mathbf{M}$, the locus of $z \in \mathbf{Z}(\mathbf{M})$ such that $zm \in \mathbf{M}^{\text{reg}/G}$, is open and Zariski dense. Similarly, the action of $\mathbf{Z}(\mathbf{M})$ on $\mathbf{M}$ induces an action of $\mathbf{Z}(\mathbf{M})$ on $\mathbf{c}_M$, and for every $m \in \mathbf{c}_M$, the locus of $z \in \mathbf{Z}(\mathbf{M})$ such that $z(m) \in \mathbf{c}_M^{\text{reg}/G}$, is open and Zariski dense. Thus, for every open subgroup $U \subset Z(M)$ we have $U(c_M^{\text{reg}/G}) = c_M$.

Note that $Z(M)$ acts smoothly on $\mathcal{H}(M)$ and induces a smooth action on $\mathcal{H}(M)_M$. Fix $h \neq 0$, and let $U \subset Z(M)$ be the stabilizer of h . Since $U(c_M^{\text{reg}/G}) = c_M$, we conclude that $h|_{c_M^{\text{reg}/G}} \neq 0$, thus $\text{res}(h) \neq 0$. $\square$

Now we show that the normalized induction map does not depend on $\mathbf{P}$.

Corollary 6.6. (a) *The normalized restriction map $r_{P;M}^G : \mathcal{H}(G)_G \rightarrow \mathcal{H}(M)_M$ and the normalized induction map $i_{P;M}^G : \hat{C}^M(M) \rightarrow \hat{C}^G(G)$ do not depend on $\mathbf{P}$.*

(b) *For every admissible representation ρ of M , the set of composition factors of $i_{P;M}^G(\rho)$ does not depend on $\mathbf{P}$.*

Proof. (a) It suffices to show the assertion for $r_{P;M}^G$. By Lemma B.4, it suffices to show that the composition $\text{res} \circ r_{P;M}^G : \mathcal{H}(G)_G \rightarrow \mathcal{H}(M)_M^{\text{reg}/G}$ does not depend on $\mathbf{P}$. On the other hand, by Corollary 6.4 for $\mathbf{H} = \mathbf{M}$, this composition equals $|\Delta_{M,G}|^{-1/2} \cdot R_M^G$.

(b) It is enough to show that the character of $\underline{i}_{P,M}^G(\rho)$ does not depend on $\mathbf{P}$. But this follows from (a) and Corollary 6.3. $\square$

6.7. Notation. By Corollary 6.6(a), we can now denote $r_{P,M}^G$ and $i_{P,M}^G$, simply by r_M^G and i_M^G , respectively.

6.8. (Stable) orbital integrals. Let $\mathbf{G}^{\text{rss}} \subset \mathbf{G}$ be the set of regular semisimple elements.

(a) In the situation of 5.4, let $\mathbf{H} = \mathbf{S} \subset \mathbf{G}$ be a maximal torus defined over F . In this case, $\mathbf{S}_{\mathbf{S}}^{\text{reg}/\mathbf{G}}$ equals $\mathbf{S}_{\mathbf{G}}^{\text{rss}} := \mathbf{S} \cap \mathbf{G}^{\text{rss}}$, and R_S^G is the map $\mathcal{H}(G) \rightarrow \mathcal{M}^\infty(S_G^{\text{rss}})$. We denote by $O_{S;G} : \mathcal{M}^\infty(S_G^{\text{rss}})^* \rightarrow \widehat{C}^G(G)$ the dual of R_S^G and call it the *orbital integral* map.

(b) Explicitly, let $\text{pr} : (\mathbf{G}/\mathbf{S}) \times \mathbf{S}_{\mathbf{G}}^{\text{rss}} \rightarrow \mathbf{S}_{\mathbf{G}}^{\text{rss}}$ be the projection. Since $\mathbf{S}$ acts on $\mathbf{S}_{\mathbf{G}}^{\text{rss}}$ trivially, it follows from 3.7(d) that R_S^G equals the composition

$$\mathcal{H}(G)_G \xrightarrow{a_{S,G}^*} \mathcal{H}([(G/S) \times S_G^{\text{rss}}]/S_G^{\text{rss}})_G \xrightarrow{\text{pr}_!} \mathcal{M}^\infty(S_G^{\text{rss}}).$$

(c) Denote by $(R_G^S)^{\text{st}} : \mathcal{H}(G)_G \rightarrow \mathcal{M}^\infty(S_G^{\text{rss}})$ the composition

$$\mathcal{H}(G)_G \xrightarrow{a_{S,G}^*} \mathcal{H}[(\mathbf{G}/\mathbf{S})(F) \times S_G^{\text{rss}}]/S_G^{\text{rss}})_G \xrightarrow{\text{pr}_!} \mathcal{M}^\infty(S_G^{\text{rss}}),$$

and let $O_{S;G}^{\text{st}} : \mathcal{M}^\infty(S_G^{\text{rss}})^* \rightarrow \widehat{C}^G(G)$ be the dual map, called the *stable orbital integral*.

6.9. Comparison. (a) Let $[g_1], \dots, [g_n] \in (\mathbf{G}/\mathbf{S})(F)$ be a set of representatives of the set of G -orbits. For every j , let $\mathbf{S}_j \subset \mathbf{G}$ be the stabilizer of $[g_j]$. Then $\mathbf{S}_j$ is a maximal torus of $\mathbf{G}$, and there is a canonical isomorphism $i_j : \mathbf{S} \xrightarrow{\sim} \mathbf{S}_j$. Explicitly, if $g_i \in \mathbf{G}(\overline{F})$ is any representative of $[g_i]$, then i_j is the map $s \mapsto g_j s g_j^{-1}$. By construction, we have the equality $O_{S;G}^{\text{st}} = \sum_{j=1}^n O_{S_j;G} \circ (i_j)_*$.

(b) Fix a Haar measure $|\omega_S|$ on S . Then for every $\gamma \in S_G^{\text{rss}}$, we can consider a “ δ -function” $\delta_\gamma \in \mathcal{M}^\infty(S_G^{\text{rss}})^*$, defined by the formula $\delta_\gamma(f|\omega_S|) = f(\gamma)$ for every $f \in C^\infty(S_G^{\text{rss}})$. Then the construction 6.8 gives us “classical” (stable) orbital integrals $O_{\gamma,G} := O_{S;G}(\delta_\gamma) \in \widehat{C}^G(G)$ (resp. $O_{\gamma,G}^{\text{st}} := O_{S;G}^{\text{st}}(\delta_\gamma) \in \widehat{C}^G(G)$). For example, observation (a) implies that a stable orbital integral is a sum of orbital integrals.

6.10. Application. Let $U \subset S_G^{\text{rss}}$ be a dense subset, and let $h \in \mathcal{H}(G)_G$ be such that $O_{\gamma,G}(h) = 0$ for every $\gamma \in U$. Then $O_{\gamma,G}(h) = 0$ for every $\gamma \in S_G^{\text{rss}}$.

Indeed, let $f \in C^\infty(S_G^{\text{rss}})$ such that $R_G^S(h) = f|\omega_S|$ (see 6.9(b)). Then for every $\gamma \in S_G^{\text{rss}}$ we have $O_{\gamma,G}(h) = f(\gamma)$. Hence, by assumption, we have $f(\gamma) = 0$ for all $\gamma \in U$. Thus $f = 0$, since f is locally constant and $U \subset S_G^{\text{rss}}$ is dense.

Corollary 6.11. *In the situation of 6.8, the following diagram is commutative*

$$\begin{array}{ccc} \mathcal{M}^\infty(S_G^{\text{rss}})^* & \xrightarrow{O_{S,G}^{\text{st}}} & \widehat{C}^G(G) \\ (|\Delta_{M,G}|^{1/2} \cdot \text{res})^* \downarrow & & \uparrow i_M^G \\ \mathcal{M}^\infty(S_M^{\text{rss}})^* & \xrightarrow{O_{S,M}^{\text{st}}} & \widehat{C}^M(M), \end{array}$$

and similarly for $O_{S,G}$ and $O_{S,M}$.

Proof. The assertion for orbital integrals is simply the dual of Corollary 6.4 for $\mathbf{H} = \mathbf{S}$. The assertion for stable orbital integrals follows from that for orbital integrals, by the observation 6.9(a) and the equality $(\mathbf{G}/\mathbf{M})(F) = G/M$. $\square$

6.12. Stable generalized functions and representations.

(a) We denote by $\widehat{C}^{\text{st}}(G) \subset \widehat{C}^G(G)$ the closure of the image of

$$O_G^{\text{st}} := \oplus_{\mathbf{S} \subset \mathbf{G}} O_{S,G}^{\text{st}} : \oplus_{\mathbf{S} \subset \mathbf{G}} \mathcal{M}^\infty(S_G^{\text{rss}})^* \rightarrow \widehat{C}^G(G),$$

where the sum is taken over the set of all maximal tori $\mathbf{S} \subset \mathbf{G}$ defined over F . Elements of $\widehat{C}^{\text{st}}(G)$ are called *stable generalized functions*.

(b) An admissible representation π of G is called *stable*, if its character χ_π is stable.

Corollary 6.13. (a) *The induction map $i_M^G : \widehat{C}^M(M) \rightarrow \widehat{C}^G(G)$ sends stable generalized functions into stable ones.*

(b) *The induction functor $i_{P,M}^G$ sends stable representations into stable ones.*

Proof. (a) Since $S_G^{\text{rss}} \subset S_M^{\text{rss}}$ is dense, the restriction map $\text{res} : \mathcal{M}^\infty(S_M^{\text{rss}}) \rightarrow \mathcal{M}^\infty(S_G^{\text{rss}})$ is injective. Thus the dual map res^* is surjective. Hence, by Corollary 6.11, we get an inclusion $i_M^G(\text{Im } O_{S,M}^{\text{st}}) \subset \widehat{C}^{\text{st}}(G)$ for every $\mathbf{S} \subset \mathbf{G}$. Since i_M^G is continuous, the assertion follows.

(b) Follows from (a) and Corollary 6.3. $\square$

APPENDIX A. A GENERALIZATION OF A THEOREM OF LUSZTIG–SPALTENSTEIN

A.1. Notation. Let F be an infinite field. All algebraic varieties and all morphisms of algebraic varieties are over F .

(a) Let $\mathbf{G}$ be a connected reductive group, $\mathbf{P} \subset \mathbf{G}$ a parabolic subgroup, $\mathbf{U} \subset \mathbf{P}$ the unipotent radical, and $\mathbf{M} \subset \mathbf{P}$ a Levi subgroup.

(b) For an algebraic variety $\mathbf{X}$, we denote the set $\mathbf{X}(F)$ by X . In particular, we have $G = \mathbf{G}(F)$, $\widetilde{G}_P = \widetilde{\mathbf{G}}_{\mathbf{P}}(F)$, etc (compare 3.8).

(c) For an $\text{Ad } P$ -invariant subset $D \subset P$, we set $\text{Ind}_P^G(D) := G \overset{P, \text{Ad}}{\times} D \subset \widetilde{G}_P$.

(d) For an $\text{Ad } M$ -invariant subset $C \subset M$, we set $\text{Ind}_M^G(C) := G \overset{M, \text{Ad}}{\times} C \subset \tilde{G}_M$, $C_P := U \cdot C \subset P$, $\text{Ind}_P^G(C_P) \subset \tilde{G}_P$ and $C_{P;G} := a_{P,G}(\text{Ind}_P^G(C_P)) \subset G$, where $a_{P,G} : \tilde{G}_P \rightarrow G$ was defined in 1.4.

From now on, we assume that $C \subset M$ is a unipotent M -conjugacy class.

A.2. Question. Does the set $C_{P;G}$ depend on the choice of $\mathbf{P} \supset \mathbf{M}$?

A.3. Remarks. (a) $C_{P;G}$ is a union of unipotent conjugacy classes in G .

(b) Let F be algebraically closed. By a theorem of Chevalley, $C_{P;G} \subset G$ is a constructible set, whose Zariski closure $\overline{C}_{P;G}$ is irreducible. This case was considered by Lusztig and Spaltenstein in [LS], and they showed that $\overline{C}_{P;G}$ does not depend on $\mathbf{P}$, using representation theory. A simpler proof of this fact was given later by Lusztig [Lu, Lem 10.3(a)].

The goal of this appendix is to generalize the result of [LS] to other fields.

A.4. Saturation. Let $\mathbf{X}$ be an algebraic variety over F , and let $A \subset X$ be a subset.

(a) We denote by $\text{sat}'(A) = \text{sat}'_{\mathbf{X}}(A) \subset X$ the union $\cup_{(\mathbf{V}, x, f)} f(x)$, taken over triples $(\mathbf{V}, x, f)$, where $\mathbf{V} \subset \mathbb{A}^1$ is an open subvariety, $x \in V$, $\mathbf{V}' := \mathbf{V} \setminus \{x\}$, and $f : \mathbf{V} \rightarrow \mathbf{X}$ is a morphism such that $f(V') \subset A$.

(b) We say that a subset $A \subset X$ is *saturated*, if $\text{sat}'(A) = A$.

(c) Let $\text{sat}(A) \subset X$ be the smallest saturated subset, containing A .

Theorem A.5. *The saturation $\text{sat}(C_{P;G})$ does not depend on $\mathbf{P}$.*

A.6. Remarks. (a) The notion of saturation is only reasonable, if the variety $\mathbf{X}$ is rationally connected.

(b) For every closed subvariety $\mathbf{Y} \subset \mathbf{X}$, the subset $\mathbf{Y}(F) \subset X$ is saturated. Also, if F is a local field, then every closed subset of X is saturated.

(c) If $\mathbf{X} = \mathbb{A}^1$, then a subset $A \subset X$ is saturated if and only if either $A = X$ or $X \setminus A$ is infinite.

(d) By (c), saturated subsets of X are not closed under finite unions. Therefore the set X does not have a topology, whose closed subsets are saturated subsets. On the other hand, our proof of Theorem A.5 indicates that in some respects saturated sets behave like closed subsets in some topology.

Lemma A.7. *Let $\mathbf{X}$ and $\mathbf{Y}$ be algebraic varieties.*

(a) *For a morphism $f : \mathbf{X} \rightarrow \mathbf{Y}$ and a subset $A \subset X$, we have an inclusion $f(\text{sat}'(A)) \subset \text{sat}'(f(A))$.*

(b) *For every $A \subset X$ and $B \subset Y$, we have $\text{sat}'(A \times B) = \text{sat}'(A) \times \text{sat}'(B)$.*

(c) *Let $\mathbf{H}$ be an algebraic group, and let $f : \mathbf{X} \rightarrow \mathbf{Y}$ be a principal $\mathbf{H}$ -bundle, locally trivial in the Zariski topology. Then for every subset $A \subset Y$ we have the equality $\text{sat}'(f^{-1}(A)) = f^{-1}(\text{sat}'(A))$.*

(d) For an $\text{Ad } P$ -invariant subset $A \subset P$, the corresponding subset $\text{Ind}_P^G(A) \subset \text{Ind}_P^G(X)$ satisfies $\text{sat}'(\text{Ind}_P^G(A)) = \text{Ind}_P^G(\text{sat}'(A))$.

(e) Let $\mathbf{Y} \subset \mathbf{X} = \mathbb{A}^n$ be an open dense subvariety. Then $\text{sat}'_X(Y) = X$.

Proof. (a) is clear.

(b) The inclusion $\subset$ follows from (a). Conversely, assume that $a \in \text{sat}'(A)$ and $b \in \text{sat}'(B)$ are defined using triples $(\mathbf{V}_a, x_a, f_a)$ and $(\mathbf{V}_b, x_b, f_b)$, respectively, where $\mathbf{V}_a$ and $\mathbf{V}_b$ are open subsets of $\mathbb{A}^1$. Then we can assume that $\mathbf{V}_a = \mathbf{V}_b \subset \mathbb{A}^1$ and $x_a = x_b$, which implies that $(a, b) \in \text{sat}'(A \times B)$.

(c) Since the saturation sat' is local in the Zariski topology, we can assume that $\mathbf{X} = \mathbf{Y} \times \mathbf{H}$. In this case the assertion follows from (b).

(d) Arguing as in 3.7(b), the assertion follows from (c).

(e) It suffices to show that for every $x \in \mathbb{A}^n(F)$ there exists a line $\mathbf{L} \subset \mathbb{A}^n$, defined over F , such that $x \in \mathbf{L}$ and $\mathbf{L} \cap \mathbf{Y} \neq \emptyset$. Consider the variety $\mathbb{P}_x$ of lines $\mathbf{L} \subset \mathbb{A}^n$ such that $x \in \mathbf{L}$. Since $\mathbf{Y} \subset \mathbb{A}^n$ is Zariski dense, the set of $\mathbf{L} \in \mathbb{P}_x$ such that $\mathbf{L} \cap \mathbf{Y} \neq \emptyset$, is Zariski dense. Since $\mathbb{P}_x \cong \mathbb{P}^{n-1}$, while F is infinite, the subset $\mathbb{P}_x(F) \subset \mathbb{P}_x$ is Zariski dense, and the assertion follows. $\square$

A.8. Remark. All the properties of sat' , formulated in Lemma A.7, have natural analogs for sat .

A.9. Relative saturation. Let $h : \mathbf{X} \rightarrow \mathbf{Y}$ be a morphism and $A \subset X$.

(a) Denote by $\text{sat}'(h; A) \subset \text{sat}'(h(A))$ the union $\cup_{(\mathbf{V}, x, f)} f(x)$, taken over all triples $(\mathbf{V}, x, f)$ in the definition of $\text{sat}'(h(A))$ (see A.4(a)) such that $f|_{\mathbf{V}'} : \mathbf{V}' \rightarrow \mathbf{Y}$ has a lift $\tilde{f}' : \mathbf{V}' \rightarrow \mathbf{X}$ with $\tilde{f}'(V') \subset A$.

(b) If h is proper, then $\text{sat}'(h; A) = h(\text{sat}'(A))$. Indeed, the valuative criterion implies that every pair $(f, \tilde{f}')$ as in (a) defines a unique morphism $\tilde{f} : \mathbf{V} \rightarrow \mathbf{X}$ such that $h \circ \tilde{f} = f$ and $\tilde{f}|_{\mathbf{V}'} = \tilde{f}'$.

(c) Let $\mathbf{X}' \subset \mathbf{X}$ be an open subvariety such that $A \subset X'$, and let $h' := h|_{\mathbf{X}'} : \mathbf{X}' \rightarrow \mathbf{Y}$ be the restriction. By definition, $\text{sat}'(h'; A) = \text{sat}'(h; A)$.

A.10. Notation. (a) $a_{\mathbf{M}, \mathbf{G}} : \tilde{\mathbf{G}}_{\mathbf{M}}^{\text{reg}} \rightarrow \mathbf{G}$ be the map defined in 1.4 and 1.5, let $\nu_{\mathbf{G}} : \mathbf{G} \rightarrow \mathbf{c}_{\mathbf{G}}$ be the Chevalley map (see 1.1(a)), and set $e_{\mathbf{G}} := \nu_{\mathbf{G}}(1) \in \mathbf{c}_{\mathbf{G}}$.

(b) Let $\mathbf{G}^{\text{der}} \subset \mathbf{G}$ be the derived group of $\mathbf{G}$, $\mathbf{Z}(\mathbf{M})$ the center of $\mathbf{M}$, and set $\mathbf{Z}_{\mathbf{M}} := (\mathbf{Z}(\mathbf{M}) \cap \mathbf{G}^{\text{der}})^0$. Then $\mathbf{Z}_{\mathbf{M}}$ is a split torus over F . Set $\mathbf{Z}_{\mathbf{M}}^{\text{reg}} := \mathbf{Z}_{\mathbf{M}} \cap \mathbf{M}^{\text{reg}}/\mathbf{G}$. Notice that since $\mathbf{Z}_{\mathbf{M}}$ acts on $\text{Lie } \mathbf{G}/\text{Lie } \mathbf{M}$ by a direct sum of non-trivial characters, the subset $\mathbf{Z}_{\mathbf{M}}^{\text{reg}} \subset \mathbf{Z}_{\mathbf{M}}$ is open and dense.

(c) Set $C_Z^{\text{reg}} := C \cdot Z_M^{\text{reg}} \subset M$. Since $C \subset M$ consists of unipotent elements, and $Z_M \subset Z(M)$, we have $C_Z^{\text{reg}} \subset M^{\text{reg}}/G$. Also C_Z^{reg} is $\text{Ad } M$ -invariant, so we can form a subset $\text{Ind}_M^G(C_Z^{\text{reg}}) \subset \tilde{\mathbf{G}}_M^{\text{reg}}$.

(d) Set $C_{P,Z}^{\text{reg}} := p^{-1}(C_Z^{\text{reg}}) = C_P \cdot Z_M^{\text{reg}} \subset P^{\text{reg}}/G$ (see Lemma 1.9(d)).

A.11. *Proof of Theorem A.5.* Consider the subset $D_P := a_{P,G}(\text{sat}'(\text{Ind}_P^G(C_P)))$ of G . Since $C_{P,G} = a_{P,G}(\text{Ind}_P^G(C_P))$, we have inclusions $C_{P,G} \subset D_P \subset \text{sat}(C_{P,G})$ (see Lemma A.7(a)), thus $\text{sat}(D_P) = \text{sat}(C_{P,G})$. It suffices to show that D_P does not depend on $\mathbf{P}$. But this follows from the following description of D_P . $\square$

Claim A.12. *We have the equality $D_P = \text{sat}'(a_{M,G}; \text{Ind}_M^G(C_Z^{\text{reg}})) \cap \nu_G^{-1}(e_G)$.*

Proof. Recall (see 1.5(d)) that morphism $a_{M,G}$ factors as $\tilde{G}_M^{\text{reg}} \xrightarrow{a_{M,P}} \tilde{G}_P \xrightarrow{a_{P,G}} G$. Notice that $a_{M,P} : \tilde{G}_M^{\text{reg}} \rightarrow \tilde{G}_P$ is an open embedding (use Lemma 1.9(e)) and it satisfies $a_{M,P}(\text{Ind}_M^G(C_Z^{\text{reg}})) = \text{Ind}_P^G(C_{P,Z}^{\text{reg}})$. Therefore, by A.9(c), we have the equality

$$\text{sat}'(a_{M,G}; \text{Ind}_M^G(C_Z^{\text{reg}})) = \text{sat}'(a_{P,G}; \text{Ind}_P^G(C_{P,Z}^{\text{reg}})).$$

Next, since $a_{P,G}$ is proper, we conclude from A.9(b) that

$$\text{sat}'(a_{P,G}; \text{Ind}_P^G(C_{P,Z}^{\text{reg}})) = a_{P,G}(\text{sat}'(\text{Ind}_P^G(C_{P,Z}^{\text{reg}}))).$$

Thus, it suffices to show the equality

$$a_{P,G}(\text{sat}'(\text{Ind}_P^G(C_P))) = a_{P,G}(\text{sat}'(\text{Ind}_P^G(C_{P,Z}^{\text{reg}}))) \cap \nu_G^{-1}(e_G).$$

Using the commutative diagram from 1.5(b) for $\mathbf{H} = \mathbf{P}$ and equality $\pi_{P,G}^{-1}(e_G) = e_P$ (see Lemma 1.9(f)), we conclude that $a_{P,G}(A) \cap \nu_G^{-1}(e_G) = a_{P,G}(A \cap \nu_{\tilde{G}_P}^{-1}(e_P))$ for every subset $A \subset \tilde{G}_P$. Thus, it suffices to show the equality

$$\text{sat}'(\text{Ind}_P^G(C_{P,Z}^{\text{reg}})) \cap \nu_{\tilde{G}_P}^{-1}(e_P) = \text{sat}'(\text{Ind}_P^G(C_P)) \subset \tilde{G}_P.$$

Using Lemma A.7(d), it suffices to show the equality

$$(A.1) \quad \text{sat}'(C_{P,Z}^{\text{reg}}) \cap \nu_P^{-1}(e_P) = \text{sat}'(C_P) \subset P.$$

Set $\mathbf{P}_{\text{un}} := \nu_P^{-1}(e_P) \subset \mathbf{P}$, and $\mathbf{P}_{\mathbf{Z}_M} := \nu_P^{-1}(\nu_P(\mathbf{Z}_M)) \subset \mathbf{P}$. Since the map $\nu_P|_{\mathbf{Z}_M} : \mathbf{Z}_M \rightarrow \mathbf{c}_P \cong \mathbf{c}_M$ is a closed embedding, the multiplication map induces an isomorphism $\mathbf{P}_{\text{un}} \times \mathbf{Z}_M \xrightarrow{\sim} \mathbf{P}_{\mathbf{Z}_M}$. Moreover, it induces a bijection $C_P \times Z_M^{\text{reg}} \xrightarrow{\sim} C_{P,Z}^{\text{reg}}$. Thus, formula (A.1) follows from the equality

$$\text{sat}'_{\mathbf{P}_{\text{un}} \times \mathbf{Z}_M}(C_P \times Z_M^{\text{reg}}) = \text{sat}'_{\mathbf{P}_{\text{un}}}(C_P) \times \text{sat}'_{\mathbf{Z}_M}(Z_M^{\text{reg}}) = \text{sat}'_{\mathbf{P}_{\text{un}}}(C_P) \times Z_M,$$

which follows from Lemma A.7(b),(e). $\square$

Corollary A.13. (a) *If F is algebraically closed, then the closure $\text{cl}(C_{P,G}) \subset G$ of $C_{P,G}$ in the Zariski topology does not depend on $\mathbf{P}$.*

(b) *If F is a local field, then the closure $\text{cl}(C_{P,G}) \subset G$ of $C_{P,G}$ in the analytic topology does not depend on $\mathbf{P}$.*

Proof. In both cases, every closed subset in G is saturated. Therefore we have inclusions $C_{P;G} \subset \text{sat}(C_{P;G}) \subset \text{cl}(C_{P;G})$, which imply that $\text{cl}(C_{P;G}) = \text{cl}(\text{sat}(C_{P;G}))$. Thus the assertion follows from Theorem A.5. $\square$

A.14. Notation. For an $\text{Ad } G$ -invariant subset $D \subset G$, we denote by $D^\heartsuit \subset D$ the union of G -conjugacy classes that are Zariski dense in (the Zariski closure of) D .

A.15. Question. Is it true that $C_{P;G}^\heartsuit$ is independent of $\mathbf{P}$?

A.16. Remarks. (a) Let F be algebraically closed. Since the number of unipotent conjugacy classes in G is finite, we conclude that $C_{P;G}^\heartsuit$ is a single conjugacy class.

(b) Let F be general. Then, by (a), $C_{P;G}^\heartsuit$ is a union of unipotent conjugacy classes, belonging to a single conjugacy class over $\overline{F}$.

Lemma A.17. *Let F be either algebraically closed or local. Then for every $\text{Ad } G$ -invariant subset $D \subset G$, we have $D^\heartsuit = \text{sat}(D)^\heartsuit$.*

Proof. Let $\text{cl}(D) \subset G$ be the closure of D in the Zariski topology if F is algebraically closed, and in the analytic topology if F is local. Then, as in the proof of Corollary A.13, we have $D \subset \text{sat}(D) \subset \text{cl}(D)$. Thus, it suffices to show that $D^\heartsuit = \text{cl}(D)^\heartsuit$.

Let $O \subset \text{cl}(D)$ be a Zariski dense G -conjugacy class, and let $\mathbf{D} \subset \mathbf{G}$ be the Zariski closure of D . Choose $x \in O$. Then the morphism $\mathbf{G} \rightarrow \mathbf{D} : g \mapsto gxg^{-1}$ is dominant. Therefore, in both cases, the corresponding map $G \rightarrow \text{cl}(D)$ is open. Thus $O \subset \text{cl}(D)$ is open, hence $O \subset D$. $\square$

Corollary A.18. *Let F be either algebraically closed or local. Then the subset $C_{P;G}^\heartsuit \subset G$ does not depend on $\mathbf{P}$.*

Proof. This follows immediately from Theorem A.5 and Lemma A.17. $\square$

A.19. Remark. We do not expect that the conclusion Lemma A.17 holds for an arbitrary field F . We wonder whether the equality $\text{sat}(C_{P;G})^\heartsuit = C_{P;G}^\heartsuit$ always holds.

APPENDIX B. ON A THEOREM OF HARISH-CHANDRA

The goal of this section is to explain the proof of the following result, usually attributed to Harish-Chandra.

Theorem B.1. *Let F be a local non-archimedean field of characteristic zero, and let $h \in \mathcal{H}(G)_G$ be such that $O_\gamma(h) = 0$ for every $\gamma \in G^{\text{rss}}$. Then $h = 0$.*

B.2. The Lie algebra analog. Let $\mathfrak{g}$ be the Lie algebra of $\mathbf{G}$, equipped with the adjoint action of G , and let $h \in \mathcal{H}(\mathfrak{g})_G$ be such that $O_x(h) = 0$ for every $x \in \mathfrak{g}^{\text{rss}}$. Then the original theorem of Harish-Chandra ([HC, Thm 3.1]) asserts that $h = 0$.

The goal of this section is to deduce Theorem B.1 from its Lie algebra analog.

B.3. G -domains. (a) Let X be a smooth analytic variety over F equipped with an action of G , let $\mathcal{H}(X)$ be the space of locally constant measures with compact support (see 3.4) and let $\mathcal{H}(X)_G$ be the space of G -coinvariants.

(b) By a G -domain in X we mean an open and closed G -invariant subset $U \subset X$. Then $\mathcal{H}(U) \subset \mathcal{H}(X)$ is a G -invariant subspace, and the map $h \mapsto 1_U \cdot h$ is a G -equivariant projection $\mathcal{H}(X) \rightarrow \mathcal{H}(U)$. Taking G -coinvariants, we get an inclusion $\mathcal{H}(U)_G \hookrightarrow \mathcal{H}(X)_G$ and a projection $\mathcal{H}(X)_G \rightarrow \mathcal{H}(U)_G \subset \mathcal{H}(X)_G : h \mapsto h|_U$.

Lemma B.4. *Let $f : X \rightarrow Y$ be a proper, surjective G -equivariant local isomorphism between smooth analytic varieties. Then the pullback map $f^* : \mathcal{H}(Y)_G \rightarrow \mathcal{H}(X)_G$ (see 3.5(b)) is injective.*

Proof. For every $m \in \mathbb{N}$, we denote by $Y_m \subset Y$ the set of all $y \in Y$ such that the cardinality of $f^{-1}(y)$ is m . The assumptions on f imply that every $Y_m \subset Y$ is a G -domain, and that Y is the disjoint union of the Y_m 's. Then every $X_m := f^{-1}(Y_m) \subset X$ is a G -domain as well, and it suffices to show that the induced map $f^* : \mathcal{H}(Y_m)_G \rightarrow \mathcal{H}(X_m)_G$ is injective. Since for every $h \in \mathcal{H}(Y_m)$ we have $f_! f^*(h) = mh$, we are done. $\square$

Proof of Theorem B.1. We carry out the proof in five steps.

Step 1. There exists a G -domain $U \ni 1$ in G such that $h|_U = 0$.

Proof. Observe first that there exist G -domains $\mathfrak{u} \ni 0$ in $\mathfrak{g}$ and $U \ni 1$ in G such that the exponential map induces an $\text{Ad } G$ -equivariant analytic isomorphism $\epsilon : \mathfrak{u} \xrightarrow{\sim} U$. Namely, if $\mathbf{G} = \mathbf{GL}_n$ the assertion is straightforward, and the general case follows from it.

We claim that this U satisfies the required property. Indeed, consider the pullback $h' := \epsilon^*(h|_U) \in \mathcal{H}(\mathfrak{u})_G \subset \mathcal{H}(\mathfrak{g})_G$. It suffices to show that $h' = 0$. For $x \in \mathfrak{g}^{\text{rss}}$ we have $O_x(h') = 0$ if $x \notin \mathfrak{u}$, since $h' \in \mathcal{H}(\mathfrak{u})_G$, and $O_x(h') = O_{\epsilon(x)}(h) = 0$ if $x \in \mathfrak{u}$ by our assumption on h . Then $h' = 0$ by [HC, Thm 3.1] (see B.2). $\square$

Step 2. For every $s \in Z(G)$ there exists a G -domain $U \ni s$ in G such that $h|_U = 0$.

Proof. Since the map $g \mapsto gs : G \rightarrow G$ is $\text{Ad } G$ -equivariant, the assertion follows from the $s = 1$ case shown in Step 1. More precisely, if $U \ni 1$ is the G -domain constructed in Step 1, then $sU \ni s$ is the G -domain such that $h|_U = 0$. $\square$

Step 3. For every semisimple $s \in G \setminus Z(G)$ there exists a G -domain $U \ni s$ in G such that $h|_U = 0$.

Proof. Let $\mathbf{H} := \mathbf{G}_s^0$ be the connected centralizer. Then $\mathbf{H} \subsetneq \mathbf{G}$, and by induction, we may assume that Theorem B.1 is valid for H .

Let $\mathbf{H}^{\text{reg}/\mathbf{G}} \subset \mathbf{H}$ and $\mathbf{c}_{\mathbf{H}}^{\text{reg}/\mathbf{G}} \subset \mathbf{c}_{\mathbf{H}}$ be the open subschemes defined in 1.2(b), let $\nu_{\mathbf{H}} : \mathbf{H}^{\text{reg}/\mathbf{G}} \rightarrow \mathbf{c}_{\mathbf{H}}^{\text{reg}/\mathbf{G}}$ be the Chevalley map (see 1.1(a)), and we denote by $\nu_H : H^{\text{reg}/G} \rightarrow \mathbf{c}_H^{\text{reg}/G}$ the induced map on F -points (compare 3.8). Note that $\mathbf{H} \subset \mathbf{G}$ is an equal rank subgroup, and $s \in H^{\text{reg}/G}$ (see 1.6).

Choose an open and closed neighbourhood $V \subset \mathbf{c}_H^{\text{reg}/G}$ of $\nu_H(s)$, and set $U' := \nu_H^{-1}(V) \subset H^{\text{reg}/G} \subset H$. Then $U' \subset H$ is a H -domain, so we can form the induced space $\text{Ind}_H^G(U')$ (see 3.7) and the G -equivariant morphism $a_{H,G} : \text{Ind}_H^G(U') \rightarrow G : [g, x] \mapsto gxg^{-1}$ (compare 1.4(c)).

Then $a_{H,G}$ is a local isomorphism because $a_{\mathbf{H},\mathbf{G}} : \text{Ind}_{\mathbf{H}}^{\mathbf{G}}(\mathbf{H}^{\text{reg}/\mathbf{G}}) \rightarrow \mathbf{G}$ is étale (see Corollary 2.6), and $a_{H,G}$ is proper because both maps $\pi_{\mathbf{H},\mathbf{G}} : \mathbf{c}_{\mathbf{H}} \rightarrow \mathbf{c}_{\mathbf{G}}$ (see 1.1(b)) $\iota_{\mathbf{H},\mathbf{G}} : \text{Ind}_{\mathbf{H}}^{\mathbf{G}}(\mathbf{H}^{\text{reg}/\mathbf{G}}) \rightarrow \mathbf{G} \times_{\mathbf{c}_{\mathbf{G}}} \mathbf{c}_{\mathbf{H}}^{\text{reg}/\mathbf{G}}$ (see Corollary 1.11) are finite. Therefore $U := \text{Im } f$ is a G -domain containing s , and we claim that $h|_U = 0$.

By Lemma B.4, the induced map $a_{H,G}^* : \mathcal{H}(U)_G \rightarrow \mathcal{H}(\text{Ind}_H^G(U'))_G$ is injective. Let $\phi : \mathcal{H}(U)_G \rightarrow \mathcal{H}(U')_H$ be the composition of $a_{H,G}^*$ and the isomorphism $\varphi_H^G : \mathcal{H}(\text{Ind}_H^G(U'))_G \xrightarrow{\sim} \mathcal{H}(U')_H$ from 3.7(c). Then ϕ is injective, it remains to show that $h' := \phi(h|_U) \in \mathcal{H}(U')_H \subset \mathcal{H}(H)_H$ is zero.

Since Theorem B.1 is valid for H , it suffices to show that $0_\gamma(h') = 0$ for all $\gamma \in H^{\text{rss}}$. This is clear for $\gamma \notin U'$, since $h' \in \mathcal{H}(U')_H$. Next, for every $\gamma \in U' \cap G^{\text{rss}}$ we have $O_\gamma(h') = O_\gamma(h) = 0$ by the assumption on h . This shows that $0_\gamma(h') = 0$ for all $\gamma \in H^{G-\text{rss}} := H \cap G^{\text{rss}}$.

Finally, since $S \cap H^{G-\text{rss}} \subset S \cap H^{\text{rss}}$ is dense for every maximal torus $\mathbf{S} \subset \mathbf{H}$, the equality $0_\gamma(h') = 0$ for every $\gamma \in H^{\text{rss}}$ now follows from 6.10. $\square$

Step 4. Let $U \subset G$ be a G -domain, and let $g = su$ be the Jordan decomposition of $g \in G$. Then $g \in U$ if and only if $s \in U$.

Proof. Set $H := G_s^0$. It suffices to show the closure of the $\text{Ad } H$ -orbit of u contains 1, hence the closure of the $\text{Ad } G$ -orbit of g contains s .

Since u is a unipotent element of H , the Zariski closure of the $\text{Ad } \mathbf{H}$ -orbit of u contains 1. Hence the assertion follows from a theorem of Kempf [Ke, Cor. 4.3]. $\square$

Step 5: Completion of the proof. By Steps 3 and 4, for every $g \in G$ there exists a G -domain $U \ni g$ in G such that $h|_U = 0$. From this the assertion follows. Indeed, choose a lift $\tilde{h} \in \mathcal{H}(G)$ of h , and let $K \subset G$ be the support of $\tilde{h}$. Since K is compact, there is a finite collection of G -domains $U_i, i = 1, \dots, n$ such that $K \subset \cup_i U_i$ and each $h_i := h|_{U_i}$ is zero. Moreover, replacing U_j by $U_j \setminus (\cup_{i=1}^{j-1} U_i)$ we can assume that the U_i 's are disjoint. Then $h = \sum_{i=1}^n h_i = 0$. $\square$

REFERENCES

- [BDK] J. Bernstein, P. Deligne and D. Kazhdan, *Trace Paley–Wiener theorem for reductive p -adic groups*, J. Analyse Math. **47** (1986) 180-192.
- [Bo] A. Borel, *Linear algebraic groups*, Second edition, Graduate Texts in Mathematics **126**, Springer-Verlag, New York, 1991.
- [HC] Harish-Chandra, *Admissible invariant distributions on reductive p -adic groups*, preface and notes by Stephen DeBacker and Paul J. Sally, Jr, University Lecture Series, **16**, American Mathematical Society, Providence, RI, 1999.
- [Ka] D. Kazhdan, *Representation of groups over close local fields*, J. Analyse Math. **47** (1986) 175-179.
- [Ke] G. Kempf, *Instability in invariant theory*, Ann. of Math. **108** (1978) 299-316.
- [LS] G. Lusztig and N. Spaltenstein, *Induced unipotent classes*, J. London Math. Soc. (2) **19** (1979) 41-52.
- [Lu] G. Lusztig, *Character sheaves on disconnected groups, II*, Represent. Theory **8** (2004) 72-124.
- [St] R. Steinberg, *Regular elements of semisimple algebraic groups*, Inst. Hautes Études Sci. Publ. Math. **25** (1965) 49-80.
- [We] A. Weil, *Adeles and algebraic groups*, Progress in Math. **23**, Birkhäuser, Boston, Mass., 1982.

INSTITUTE OF MATHEMATICS, THE HEBREW UNIVERSITY OF JERUSALEM, GIVAT-RAM, JERUSALEM, 91904, ISRAEL

E-mail address: kazhdan@math.huji.ac.il

INSTITUTE OF MATHEMATICS, THE HEBREW UNIVERSITY OF JERUSALEM, GIVAT-RAM, JERUSALEM, 91904, ISRAEL

E-mail address: vyakov@math.huji.ac.il